arXiv:physics/0110044v1 [physics.gen-ph] 15 Oct 2001

A Piece of the Lepton Theory from a Probability

Gunn Alex Quznetsov*

February 2, 2008

Abstract

A masses of a leptons deduced from a representation of a probability density vector by a spinors. A massive W and Z bosons and a massless A boson are obtained from a transformations for which a density vector is invariant.

I use the following denotations:

$$1_2 = \left[\begin{array}{cc} 1 & 0 \\ 0 & 1 \end{array}\right], 0_2 = \left[\begin{array}{cc} 0 & 0 \\ 0 & 0 \end{array}\right]$$

and for $k \geq 2$:

$$1_{2k} = \left[\begin{array}{cc} 1_k & 0_k \\ 0_k & 1_k \end{array}\right], 0_{2k} = \left[\begin{array}{cc} 0_k & 0_k \\ 0_k & 0_k \end{array}\right].$$

$$\sigma_1 = \left[\begin{array}{cc} 0 & 1 \\ 1 & 0 \end{array}\right], \sigma_2 = \left[\begin{array}{cc} 0 & -i \\ i & 0 \end{array}\right], \sigma_3 = \left[\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array}\right]$$

are the Pauli matrices.

The Clifford pentad [2] $\overset{\circ}{\beta}$ is:

---

*quznets@yahoo.com

$$\begin{gathered}\beta_1=\begin{bmatrix}\sigma_1 & 0_2\\ 0_2 & -\sigma_1\end{bmatrix},\beta_2=\begin{bmatrix}\sigma_2 & 0_2\\ 0_2 & -\sigma_2\end{bmatrix},\beta_3=\begin{bmatrix}\sigma_3 & 0_2\\ 0_2 & -\sigma_3\end{bmatrix},\\ \gamma_0=\begin{bmatrix}0_2 & 1_2\\ 1_2 & 0_2\end{bmatrix}=\beta_5,\beta_4=i\begin{bmatrix}0_2 & 1_2\\ -1_2 & 0_2\end{bmatrix},\end{gathered}\tag{1}$$

$$\beta_0=\begin{bmatrix}1_2 & 0_2\\ 0_2 & 1_2\end{bmatrix},\ \gamma_5=\begin{bmatrix}1_2 & 0_2\\ 0_2 & -1_2\end{bmatrix}.$$

1 Masses

Let

$$\langle\rho\left(t,x,y,z\right),j_x\left(t,x,y,z\right),j_y\left(t,x,y,z\right),j_z\left(t,x,y,z\right)\rangle$$

be a probability current 3+1 vector field [3] and $\psi\left(t,x,y,z\right)$ be any complex spinor field:

$$\psi=|\psi|\begin{bmatrix}\exp\left(i\overset{*}{\gamma}\right)\cos\left(\overset{*}{\beta}\right)\cos\left(\overset{*}{\alpha}\right)\\ \exp\left(i\overset{*}{\theta}\right)\sin\left(\overset{*}{\beta}\right)\cos\left(\overset{*}{\alpha}\right)\\ \exp\left(i\overset{*}{\varphi}\right)\cos\left(\overset{*}{\chi}\right)\sin\left(\overset{*}{\alpha}\right)\\ \exp\left(i\overset{*}{\upsilon}\right)\sin\left(\overset{*}{\chi}\right)\sin\left(\overset{*}{\alpha}\right)\end{bmatrix}.$$

In this case the following system of equations:

$$\left\{\begin{array}{c}\psi^\dagger\psi=\rho,\\ \psi^\dagger\beta_1\psi=j_x,\\ \psi^\dagger\beta_2\psi=j_y,\\ \psi^\dagger\beta_3\psi=j_z\end{array}\right|\tag{2}$$

has got the following form:

$$\left\{\begin{array}{c} \psi^{\dagger}\psi=\rho, \\ |\psi|^{2}\left(\begin{array}{c} \cos^{2}\left(\overset{*}{\alpha}\right)\sin\left(2\,\overset{*}{\beta}\right)\cos\left(\overset{*}{\theta}-\overset{*}{\gamma}\right)- \\ -\sin^{2}\left(\overset{*}{\alpha}\right)\sin\left(2\,\overset{*}{\chi}\right)\cos\left(\overset{*}{\upsilon}-\overset{*}{\varphi}\right) \end{array}\right)=j_{x}, \\ |\psi|^{2}\left(\begin{array}{c} \cos^{2}\left(\overset{*}{\alpha}\right)\sin\left(2\,\overset{*}{\beta}\right)\sin\left(\overset{*}{\theta}-\overset{*}{\gamma}\right)- \\ -\sin^{2}\left(\overset{*}{\alpha}\right)\sin\left(2\,\overset{*}{\chi}\right)\sin\left(\overset{*}{\upsilon}-\overset{*}{\varphi}\right) \end{array}\right)=j_{y}, \\ |\psi|^{2}\left(\cos^{2}\left(\overset{*}{\alpha}\right)\cos\left(2\,\overset{*}{\beta}\right)-\sin^{2}\left(\overset{*}{\alpha}\right)\cos\left(2\,\overset{*}{\chi}\right)\right)=j_{z}. \end{array}\right|$$

Hence for every probability current vector $\langle\rho,j_x,j_y,j_z\rangle$: the spinor ψ, obeyed to this system, exists.

The operator $\widehat{U}\left(t,\triangle t\right)$, which acts in the set of these spinors, is denoted as the evolution operator for the spinor $\psi\left(t,x,y,z\right)$, if:

$$\psi\left(t+\triangle t,x,y,z\right)=\widehat{U}\left(t,\triangle t\right)\psi\left(t,x,y,z\right).$$

$\widehat{U}\left(t,\triangle t\right)$ is a linear operator.

The set of the spinors, for which $\widehat{U}\left(t,\triangle t\right)$ is the evolution operator, is denoted as the operator $\widehat{U}\left(t,\triangle t\right)$ space.

The operator space is the linear space.

Let for an infinitesimal $\triangle t$:

$$\widehat{U}\left(t,\triangle t\right)=1+i\triangle t\widehat{H}\left(t\right).$$

Hence for an elements of the operator $\widehat{U}\left(t,\triangle t\right)$ space:

$$i\widehat{H}=\partial_t.$$

If the functions ρ, j_x, j_y, j_z fulfill to the continuity equation [3]:

$$\partial_t\rho+\partial_x j_x+\partial_y j_y+\partial_z j_z=0$$

then:

$$\left(\left(\partial_t\psi^{\dagger}\right)\beta_0+\left(\partial_x\psi^{\dagger}\right)\beta_1+\left(\partial_y\psi^{\dagger}\right)\beta_2+\left(\partial_z\psi^{\dagger}\right)\beta_3\right)\psi=$$

$$=-\psi^{\dagger}\left(\left(\beta_0\partial_t+\beta_1\partial_x+\beta_2\partial_y+\beta_3\partial_z\right)\psi\right).$$

Let:

$$\widehat{Q}=\left(i\widehat{H}+\beta_1\partial_x+\beta_2\partial_y+\beta_3\partial_z\right).$$

Hence:

$$\psi^\dagger\widehat{Q}^\dagger\psi=-\psi^\dagger\widehat{Q}\psi.$$

Hence $i\widehat{Q}\left(t,x,y,z\right)$ is the Hermitian for the matrix product operator. Hence a real functions $\varphi_{i,j}\left(t,x,y,z\right)$ and $\varpi_{i,j}\left(t,x,y,z\right)$ for which:

$$-i\widehat{Q}=$$

$$\left[\begin{array}{cccc}
\varphi_{1,1} & \varphi_{1,2}+i\varpi_{1,2} & \varphi_{1,3}+i\varpi_{1,3} & \varphi_{1,4}+i\varpi_{1,4}\\
\varphi_{1,2}-i\varpi_{1,2} & \varphi_{2,2} & \varphi_{2,3}+i\varpi_{2,3} & \varphi_{2,4}+i\varpi_{2,4}\\
\varphi_{1,3}-i\varpi_{1,3} & \varphi_{2,3}-i\varpi_{2,3} & \varphi_{3,3} & \varphi_{3,4}+i\varpi_{3,4}\\
\varphi_{1,4}-i\varpi_{1,4} & \varphi_{2,4}-i\varpi_{2,4} & \varphi_{3,4}-i\varpi_{3,4} & \varphi_{4,4}
\end{array}\right]$$

exist.

Let G_t, G_z, K_t and K_z are the solution of the following system of equations:

$$\left\{\begin{array}{c}
G_t+G_z+K_t+K_z=\varphi_{1,1},\\
G_t-G_z+K_t-K_z=\varphi_{2,2},\\
G_t-G_z-K_t+K_z=\varphi_{3,3},\\
G_t+G_z-K_t-K_z=\varphi_{4,4};
\end{array}\right|$$

G_x and K_x are the solution of the following system of equations:

$$\left\{\begin{array}{c}
G_x+K_x=\varphi_{1,2},\\
-G_x+K_x=\varphi_{3,4};
\end{array}\right|$$

G_y and K_y are the solution of the following system of equations:

$$\left\{\begin{array}{c}
-G_y-K_y=\varpi_{1,2},\\
G_y-K_x=\varpi_{3,4}.
\end{array}\right|$$

In this case:

$$\begin{gathered}
-i\widehat{Q}=\\
=\left(G_t\beta_0+G_x\beta_1+G_y\beta_2+G_z\beta_3\right)+\\
+\left(K_t\beta_0+K_x\beta_1+K_y\beta_2+K_z\beta_3\right)\gamma_5+
\end{gathered}$$

$$+\left[\begin{array}{cccc} 0 & 0 & \varphi_{1,3}+i\varpi_{1,3} & \varphi_{1,4}+i\varpi_{1,4} \\ 0 & 0 & \varphi_{2,3}+i\varpi_{2,3} & \varphi_{2,4}+i\varpi_{2,4} \\ \varphi_{1,3}-i\varpi_{1,3} & \varphi_{2,3}-i\varpi_{2,3} & 0 & 0 \\ \varphi_{1,4}-i\varpi_{1,4} & \varphi_{2,4}-i\varpi_{2,4} & 0 & 0 \end{array}\right].$$

If

$$\left\{\begin{array}{c} (M_0+M_{z,0})=\varphi_{1,3}, \\ (M_0-M_{z,0})=\varphi_{2,4}, \end{array}\right|$$

$$\left\{\begin{array}{c} (M_4+M_{z,4})=\varpi_{1,3}, \\ (M_4-M_{z,4})=\varpi_{2,4}, \end{array}\right|$$

$$\left\{\begin{array}{c} (M_{x,0}+M_{y,4})=\varphi_{1,4}, \\ (M_{x,0}-M_{y,4})=\varphi_{2,3}, \end{array}\right|$$

$$\left\{\begin{array}{c} (M_{x,4}+M_{y,0})=\varpi_{1,4}, \\ (M_{x,4}-M_{y,0})=\varpi_{2,3} \end{array}\right|$$

then

$$\left[\begin{array}{cccc} 0 & 0 & \varphi_{1,3}+i\varpi_{1,3} & \varphi_{1,4}+i\varpi_{1,4} \\ 0 & 0 & \varphi_{2,3}+i\varpi_{2,3} & \varphi_{2,4}+i\varpi_{2,4} \\ \varphi_{1,3}-i\varpi_{1,3} & \varphi_{2,3}-i\varpi_{2,3} & 0 & 0 \\ \varphi_{1,4}-i\varpi_{1,4} & \varphi_{2,4}-i\varpi_{2,4} & 0 & 0 \end{array}\right]=$$

$$=M_0\gamma_0+M_4\beta_4-M_{x,0}\gamma_\zeta^0-M_{x,4}\zeta^4+M_{y,0}\gamma_\eta^0+M_{y,4}\eta^4-M_{z,0}\gamma_\theta^0-M_{z,4}\theta^4;$$

here γ_ζ^0, ζ^4, γ_η^0, η^4, γ_θ^0, θ^4 are the chromatic pentads [4], [5] members and γ_0 and β_4 is the light pentad $\overset{\circ}{\beta}$ members. Since in this paper I will not consider a quarks then everywhere below:

$$M_{x,0}=M_{x,4}=M_{y,0}=M_{y,4}=M_{z,0}=M_{z,4}=0$$

hence:

$$\begin{gathered}-i\widehat{Q}=\\=\left(G_t\beta_0+G_x\beta_1+G_y\beta_2+G_z\beta_3\right)+\\+\left(K_t\beta_0+K_x\beta_1+K_y\beta_2+K_z\beta_3\right)\gamma_5+\\+M_0\gamma_0+M_4\beta_4.\end{gathered}$$

$\{\beta_1,\beta_2,\beta_3,\beta_4,\gamma_0\}$ is the Clifford pentad.

If $j_x=\rho u_x$, $j_y=\rho u_y$, $j_z=\rho u_z$ then u_x, u_y, u_z are the components of the average velocity. Hence β_1, β_2, β_3 define the components of the average velocity (2).

If

$$j_{x_5}=\psi^\dagger\gamma_0\psi,\ j_{x_4}=\psi^\dagger\beta_4\psi,\ j_{x_5}=\rho u_{x_5},\ j_{x_4}=\rho u_{x_4}$$

then

$$\begin{gathered}u_{x_5}=\sin\left(2\overset{*}{\alpha}\right)\left[\begin{array}{c}\cos\left(\overset{*}{\beta}\right)\cos\left(\overset{*}{\chi}\right)\cos\left(\overset{*}{\gamma}-\overset{*}{\varphi}\right)+\\+\sin\left(\overset{*}{\beta}\right)\sin\left(\overset{*}{\chi}\right)\cos\left(\overset{*}{\theta}-\overset{*}{\upsilon}\right)\end{array}\right],\\u_{x_4}=\sin\left(2\overset{*}{\alpha}\right)\left[\begin{array}{c}\cos\left(\overset{*}{\beta}\right)\cos\left(\overset{*}{\chi}\right)\sin\left(\overset{*}{\gamma}-\overset{*}{\varphi}\right)+\\+\sin\left(\overset{*}{\beta}\right)\sin\left(\overset{*}{\chi}\right)\sin\left(\overset{*}{\theta}-\overset{*}{\upsilon}\right)\end{array}\right]\end{gathered}$$

and if $\rho\neq 0$ then

$$u_x^2+u_y^2+u_z^2+u_{x_5}^2+u_{x_4}^2=1. \tag{3}$$

From [1] the maximal velocity of the information propagation in the space-time is 1.

Hence of only all five elements of the Clifford pentad lends the entire kit of the velocity components and, for the completeness, yet two "space" coordinates x_5 and x_4 should be added to our three x,y,z.

Let

$$\begin{gathered}\mathbf{\Psi}\left(t,x,y,z,x_5,x_4\right)=\\=\psi\left(t,x,y,z\right)\exp\left(-i\left(x_5M_0\left(t,x,y,z\right)+x_4M_4\left(t,x,y,z\right)\right)\right).\end{gathered}$$

In this case the motion equation is the following:

$$\begin{gathered}\beta_0 i\partial_t \mathbf{\Psi} + \beta_1 i\partial_x \mathbf{\Psi} + \beta_2 i\partial_y \mathbf{\Psi} + \beta_3 i\partial_z \mathbf{\Psi} + \gamma_0 i\partial_{x_5} \mathbf{\Psi} + \beta_4 i\partial_{x_4} \mathbf{\Psi} + \\ + \left(G_t\beta_0 + G_x\beta_1 + G_y\beta_2 + G_z\beta_3\right)\mathbf{\Psi} + \\ + \left(K_t\beta_0 + K_x\beta_1 + K_y\beta_2 + K_z\beta_3\right)\gamma_5 \mathbf{\Psi} = 0\end{gathered} \tag{4}$$

Let a evolution operator $\widehat{U}\left(t, \triangle t\right)$ be denoted as *a Planck evolution operator* if a tiny positive real number h and a functions $N_\varphi\left(t,x,y,z\right)$ and $N_\varpi\left(t,x,y,z\right)$, having a range of values in the set of the integer numbers, exist for which:

$$M_0 = N_\varphi h \text{ and } M_4 = N_\varpi h.$$

Let $-\frac{\pi}{h} \leq x_5 \leq \frac{\pi}{h}$, $-\frac{\pi}{h} \leq x_4 \leq \frac{\pi}{h}$,
$\mathbf{\Psi}\left(t,x,y,z,\pm\frac{\pi}{h},x_4\right) = 0$ and $\mathbf{\Psi}\left(t,x,y,z,x_5,\pm\frac{\pi}{h}\right) = 0$.
In this case the Fourier series for $\mathbf{\Psi}$ is of the following form:

$$\begin{gathered}\mathbf{\Psi}\left(t,x,y,z,x_5,x_4\right) = \\ = \psi\left(t,x,y,z\right)\sum_{\nu,\kappa}\delta_{-\nu,N_\varphi(t,x,y,z)}\delta_{-\kappa,N_\varpi(t,x,y,z)}\exp\left(-ih\left(\nu x_5 + \kappa x_4\right)\right).\end{gathered}$$

Here:

$$\begin{aligned}\delta_{-\nu,N_\varphi} &= \frac{h}{2\pi}\int_{-\frac{\pi}{h}}^{\frac{\pi}{h}}\exp\left(ih\left(\nu x_5\right)\right)\exp\left(iN_\varphi h x_5\right)dx_5 = \frac{\sin\left(\pi\left(\nu + N_\varphi\right)\right)}{\pi\left(\nu + N_\varphi\right)}, \\ \delta_{-\kappa,N_\varpi} &= \frac{h}{2\pi}\int_{-\frac{\pi}{h}}^{\frac{\pi}{h}}\exp\left(ih\left(\kappa x_4\right)\right)\exp\left(iN_\varpi h x_4\right)dx_4 = \frac{\sin\left(\pi\left(\kappa + N_\varpi\right)\right)}{\pi\left(\kappa + N_\varpi\right)}.\end{aligned}$$

If denote:

$$\phi\left(t,x,y,z,-\nu,-\kappa\right) = \psi\left(t,x,y,z\right)\delta_{\nu,N_\varphi(t,x,y,z)}\delta_{\kappa,N_\varpi(t,x,y,z)}$$

then

$$\begin{gathered}\mathbf{\Psi}\left(t,x,y,z,x_5,x_4\right) = \\ = \sum_{\nu,\kappa}\phi\left(t,x,y,z,\nu,\kappa\right)\exp\left(-ih\left(\nu x_5 + \kappa x_4\right)\right).\end{gathered}$$

From the properties of δ in every point $\langle t,x,y,z\rangle$: either

$$\mathbf{\Psi}\left(t,x,y,z,x_5,x_4\right) = 0$$

or an integer numbers ν_0 and κ_0 exist for which:

$$\mathbf{\Psi}\left(t,x,y,z,x_5,x_4\right)=\phi\left(t,x,y,z,\nu_0,\kappa_0\right)\exp\left(-ih\left(\nu_0x_5+\kappa_0x_4\right)\right). \qquad (5)$$

That is for the every space-time point: either this point is empty or single mass is placed in this point.

Let on the space of these spinors the scalar product $\mathbf{\Phi}*\mathbf{\Psi}$ be denoted as the following:

$$\mathbf{\Phi}*\mathbf{\Psi}=\left(\frac{h}{2\pi}\right)^2\int_{-\frac{\pi}{h}}^{\frac{\pi}{h}}dx_5\int_{-\frac{\pi}{h}}^{\frac{\pi}{h}}dx_4\cdot\left(\mathbf{\Phi}^\dagger\mathbf{\Psi}\right).$$

In this case:

$$\mathbf{\Psi}*\beta_\mu\mathbf{\Psi}=\psi^\dagger\beta_\mu\psi.$$

for $0\leq\mu\leq 3$

Hence from (2):

$$\left\{\begin{array}{c}\mathbf{\Psi}^\dagger*\mathbf{\Psi}=\rho,\\ \mathbf{\Psi}^\dagger*\beta_1\mathbf{\Psi}=j_x,\\ \mathbf{\Psi}^\dagger*\beta_2\mathbf{\Psi}=j_y,\\ \mathbf{\Psi}^\dagger*\beta_3\mathbf{\Psi}=j_z.\end{array}\right|$$

1.1 Bi-zero-nonzero-mass state

Let

$$\begin{gathered}\mathbf{\Psi}\left(t,x,y,z,x_5,x_4\right)=\\ =\phi\left(t,x,y,z,0,0\right)+\phi\left(t,x,y,z,n,k\right)\exp\left(ih\left(nx_5+kx_4\right)\right).\end{gathered}$$

Let ϵ_μ $(1\leq k\leq 4)$ be a basis in which pentad $\overset{\circ}{\beta}$ has got a form (1) and let

$$\begin{gathered}\mathbf{\Psi}\left(x_5,x_4\right)=\\ =\sum_{r=1}^4\phi_r\left(0,0\right)\epsilon_r+\exp\left(-ih\left(nx_5+kx_4\right)\right)\sum_{k=1}^4\phi_k\left(n,k\right)\epsilon_k\end{gathered} \qquad (6)$$

Hence in the basis

$$\langle \epsilon_r, \exp\left(-ih\left(nX + kY\right)\right)\epsilon_k\rangle :$$

a 8-components bi-spinor:

$$\Psi = \begin{bmatrix} \phi_1\left(0,0\right) \\ \phi_2\left(0,0\right) \\ \phi_3\left(0,0\right) \\ \phi_4\left(0,0\right) \\ \phi_1\left(n,k\right) \\ \phi_2\left(n,k\right) \\ \phi_3\left(n,k\right) \\ \phi_4\left(n,k\right) \end{bmatrix}$$

corresponds to $\mathbf{\Psi}$.
From (5): in every point $\langle t, x, y, z\rangle$:

$$\Psi = \begin{bmatrix} \phi_1\left(0,0\right) \\ \phi_2\left(0,0\right) \\ \phi_3\left(0,0\right) \\ \phi_4\left(0,0\right) \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \text{ or } \Psi = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ \phi_1\left(n,k\right) \\ \phi_2\left(n,k\right) \\ \phi_3\left(n,k\right) \\ \phi_4\left(n,k\right) \end{bmatrix} \tag{7}$$

of δ characteristics.
Let us denote:

$$\phi_1\epsilon_1 + \phi_2\epsilon_2 = \phi_L \text{ and } \phi_3\epsilon_3 + \phi_4\epsilon_4 = \phi_R.$$

Hence from (6):

$$\begin{gathered} \mathbf{\Psi}\left(x_5, x_4\right) = \phi_L\left(0,0\right) + \phi_R\left(0,0\right) + \\ + \exp\left(-ih\left(nx_5 + kx_4\right)\right)\left(\phi_L\left(n,k\right) + \phi_R\left(n,k\right)\right). \end{gathered} \tag{8}$$

If use denotation:

$$\underline{\vartheta} = \begin{bmatrix} \vartheta & 0_4 \\ 0_4 & \vartheta \end{bmatrix},$$

$$\gamma = \left[\begin{array}{cc} -\gamma_0 & 0_4 \\ 0_4 & \gamma_0 \end{array}\right], \beta = \left[\begin{array}{cc} -\beta_4 & 0_4 \\ 0_4 & \beta_4 \end{array}\right]$$

and

$$\underline{n} = \left[\begin{array}{cc} 0_4 & 0_4 \\ 0_4 & n1_4 \end{array}\right], \underline{k} = \left[\begin{array}{cc} 0_4 & 0_4 \\ 0_4 & k1_4 \end{array}\right]$$

then the motion equation is the following:

$$\begin{gathered} \underline{\beta_0} i\partial_t \Psi + \underline{\beta_1} i\partial_x \Psi + \underline{\beta_2} i\partial_y \Psi + \underline{\beta_3} i\partial_z \Psi + \\ -h\underline{n}\gamma\Psi - h\underline{k}\beta\Psi + \\ + \left(G_t\underline{\beta_0} + G_x\underline{\beta_1} + G_y\underline{\beta_2} + G_z\underline{\beta_3}\right)\Psi + \\ + \left(K_t\underline{\beta_0} + K_x\underline{\beta_1} + K_y\underline{\beta_2} + K_z\underline{\beta_3}\right)\underline{\gamma_5}\Psi = 0 \end{gathered} \tag{9}$$

and

$$\left\{\begin{array}{c} \Psi^\dagger\Psi = \rho, \\ \Psi^\dagger\underline{\beta_1}\Psi = j_x, \\ \Psi^\dagger\underline{\beta_2}\Psi = j_y, \\ \Psi^\dagger\underline{\beta_3}\Psi = j_z. \end{array}\right|$$

If use the following denotation: $t = x_0, x = x_1, y = x_2, z = x_3, \partial_\mu = \frac{\partial}{\partial x_\mu}$ then the lagrangian has got the following form:

$$\begin{gathered} \mathcal{L}_f = 0.5i\left(\left(\sum_{\mu=0}^{3}\Psi^\dagger\underline{\beta_\mu}\partial_\mu\Psi\right) - \left(\sum_{\mu=0}^{3}\partial_\mu\Psi^\dagger\underline{\beta_\mu}\Psi\right)\right) - \\ - \left(\Psi^\dagger h\underline{n}\gamma\Psi + \Psi^\dagger h\underline{k}\beta\Psi\right) + \\ +\Psi^\dagger\left(\sum_{\mu=0}^{3}G_{x_\mu}\underline{\beta_\mu}\right)\Psi + \Psi^\dagger\left(\sum_{\mu=0}^{3}K_{x_\mu}\underline{\beta_\mu}\right)\underline{\gamma_5}\Psi. \end{gathered} \tag{10}$$

This lagrangian is invariant for the rotation of xOy, yOz, xOz and for the Lorentz transformation of tOx, tOy, tOz and G_{x_k} and K_{x_k} behaves as the 4-vector fields [6].

1.1.1 Transformations

If U is an 8×8 complex matrix, $\Psi' = U\Psi$ and

$$\left\{\begin{array}{c} \Psi'^\dagger\underline{\beta_1}\Psi' = j_x, \\ \Psi'^\dagger\underline{\beta_2}\Psi' = j_y, \\ \Psi'^\dagger\underline{\beta_3}\Psi' = j_z \end{array}\right| \tag{11}$$

then for $1 \leq k \leq 3$: $U^{\dagger}\underline{\beta_k}U = \underline{\beta_k}$. In this case a real numbers $a”, b”, c”, g”, u”, v”, k, s, a‘, b‘, c‘, g‘, u‘, v‘, k‘, s‘$ exist for which:

$$\underline{U} = \left[\begin{array}{cccc} (a” + b”i)\,1_2 & 0_2 & (c” + ig”)\,1_2 & 0_2 \\ 0_2 & (a‘ + b‘i)\,1_2 & 0_2 & (c‘ + ig‘)\,1_2 \\ (u” + iv”)\,1_2 & 0_2 & (k” + is”)\,1_2 & 0_2 \\ 0_2 & (u‘ + iv‘)\,1_2 & 0_2 & (k‘ + is‘)\,1_2 \end{array} \right].$$

If $\Psi'^{\dagger}\Psi' = \rho$ then $U^{\dagger}U = 1_8$. Hence:

$$\begin{array}{c} v”^2 + b”^2 + u”^2 + a”^2 = 1, \\ c”^2 + g”^2 + k”^2 + s”^2 = 1, \end{array}$$

$$s” = -\frac{a”g”u” - u”b”c” + a”c”v” + b”g”v”}{u”^2 + v”^2},$$

$$k” = \frac{-u”a”c” - u”b”g” + v”a”g” - b”c”v”}{u”^2 + v”^2}.$$

$$\begin{array}{c} v‘^2 + b‘^2 + u‘^2 + a‘^2 = 1, \\ c‘^2 + g‘^2 + k‘^2 + s‘^2 = 1, \end{array}$$

$$s‘ = -\frac{a‘g‘u‘ - u‘b‘c‘ + a‘c‘v‘ + b‘g‘v‘}{u‘^2 + v‘^2},$$

$$k‘ = \frac{-u‘a‘c‘ - u‘b‘g‘ + v‘a‘g‘ - b‘c‘v‘}{u‘^2 + v‘^2}.$$

$\underline{U}$ has got 4 eigenvalues: $\exp(i\alpha_1)$, $\exp(i\alpha_2)$, $\exp(i\alpha_3)$, $\exp(i\alpha_4)$ for 8 orthogonal eigenvectors:

$\varepsilon_{1,1}$, $\varepsilon_{1,2}$, $\varepsilon_{2,1}$, $\varepsilon_{2,2}$, $\varepsilon_{3,1}$, $\varepsilon_{3,2}$, $\varepsilon_{4,1}$, $\varepsilon_{4,2}$.

Let

$$K = \left[\begin{array}{cccccccc} \varepsilon_{1,1} & \varepsilon_{1,2} & \varepsilon_{2,1} & \varepsilon_{2,2} & \varepsilon_{3,1} & \varepsilon_{3,2} & \varepsilon_{4,1} & \varepsilon_{4,2} \end{array} \right].$$

Let θ_1, θ_2, θ_3, θ_4 be the solution of the system of the equations:

$$\left\{ \begin{array}{l} \theta_1+\theta_2+\theta_3+\theta_4=\alpha_1, \\ \theta_1+\theta_2-\theta_3-\theta_4=\alpha_1, \\ \theta_1-\theta_2+\theta_3-\theta_4=\alpha_1, \\ \theta_1-\theta_2-\theta_3+\theta_4=\alpha_1. \end{array} \right|$$

and

$$U_1=\exp\left(i\theta_1\right)1_8,$$

$$U_2=K\left[\begin{array}{cc} \exp\left(i\theta_2\right)1_4 & 0_4 \\ 0_4 & \exp\left(-i\theta_2\right)1_4 \end{array}\right]K^{\dagger},$$

$$U_3=K\left[\begin{array}{cccc} \exp\left(i\theta_3\right)1_2 & 0_2 & 0_2 & 0_2 \\ 0_2 & \exp\left(-i\theta_3\right)1_2 & 0_2 & 0_2 \\ 0_2 & 0_2 & \exp\left(i\theta_3\right)1_2 & 0_2 \\ 0_2 & 0_2 & 0_2 & \exp\left(-i\theta_3\right)1_2 \end{array}\right]K^{\dagger},$$

$$U_4=K\left[\begin{array}{cccc} \exp\left(i\theta_4\right)1_2 & 0_2 & 0_2 & 0_2 \\ 0_2 & \exp\left(-i\theta_4\right)1_2 & 0_2 & 0_2 \\ 0_2 & 0_2 & \exp\left(-i\theta_4\right)1_2 & 0_2 \\ 0_2 & 0_2 & 0_2 & \exp\left(i\theta_4\right)1_2 \end{array}\right]K^{\dagger}.$$

In this case:

$$U_1U_2U_3U_4=U$$

and

$$U_2=\left[\begin{array}{cccc} \exp\left(i\theta_2\right)1_2 & 0_2 & 0_2 & 0_2 \\ 0_2 & \exp\left(-i\theta_2\right)1_2 & 0_2 & 0_2 \\ 0_2 & 0_2 & \exp\left(i\theta_2\right)1_2 & 0_2 \\ 0_2 & 0_2 & 0_2 & \exp\left(-i\theta_2\right)1_2 \end{array}\right]$$

and a real number a,b,c,g,u,v,k,s exist for which:

$$U_3U_4=\left[\begin{array}{cccc} \left(a+ib\right)1_2 & 0_2 & \left(c+ig\right)1_2 & 0_2 \\ 0_2 & \left(u+iv\right)1_2 & 0_2 & \left(k+is\right)1_2 \\ \left(-c+ig\right)1_2 & 0_2 & \left(a-ib\right)1_2 & 0_2 \\ 0_2 & \left(-k+is\right)1_2 & 0_2 & \left(u-iv\right)1_2 \end{array}\right]$$

and

$$\begin{array}{c} a^2+b^2+c^2+g^2=1, \\ u^2+v^2+r^2+s^2=1. \end{array}$$

If

$$U^{(+)}=\left[\begin{array}{cccc} 1_2 & 0_2 & 0_2 & 0_2 \\ 0_2 & (u+iv)\,1_2 & 0_2 & (k+is)\,1_2 \\ 0_2 & 0_2 & 1_2 & 0_2 \\ 0_2 & (-k+is)\,1_2 & 0_2 & (u-iv)\,1_2 \end{array}\right] \tag{12}$$

and

$$U^{(-)}=\left[\begin{array}{cccc} (a+ib)\,1_2 & 0_2 & (c+ig)\,1_2 & 0_2 \\ 0_2 & 1_2 & 0_2 & 0_2 \\ (-c+ig)\,1_2 & 0_2 & (a-ib)\,1_2 & 0_2 \\ 0_2 & 0_2 & 0_2 & 1_2 \end{array}\right]$$

then

$$U_3U_4=U^{(-)}U^{(+)}=U^{(+)}U^{(-)}.$$

B-boson

$$U_1U_2=\left[\begin{array}{cccc} e^{i(\theta_1+\theta_2)} & 0 & 0 & 0 \\ 0 & e^{i(\theta_1-\theta_2)} & 0 & 0 \\ 0 & 0 & e^{i(\theta_1+\theta_2)} & 0 \\ 0 & 0 & 0 & e^{i(\theta_1-\theta_2)} \end{array}\right]$$

Let χ and ς be the solution of the following set of equations:

$$\left\{\begin{array}{c} 0.5\chi+\varsigma=\theta_1+\theta_2, \\ \chi+\varsigma=\theta_1-\theta_2, \end{array}\right|$$

i.e.:

$$\begin{array}{c} \chi=-4\theta_2, \\ \varsigma=\theta_1+3\theta_2. \end{array}$$

Let

$$\overbrace{U}=\exp\left(i\varsigma\right)1_8$$

and

$$\widetilde{U} = \begin{bmatrix} \exp\left(i\frac{\chi}{2}\right)1_2 & 0_2 & 0_2 & 0_2 \\ 0_2 & \exp\left(i\chi\right)1_2 & 0_2 & 0_2 \\ 0_2 & 0_2 & \exp\left(i\frac{\chi}{2}\right)1_2 & 0_2 \\ 0_2 & 0_2 & 0_2 & \exp\left(i\chi\right)1_2 \end{bmatrix}.$$

In that case:

$$\widetilde{U}\overbrace{U} = U_1 U_2.$$

Let g_1 be a positive real number and for $\mu \in \{t, x, y, z\}$: F_μ and B_μ be the solutions of the following systems of the equations:

$$\left\{ \begin{array}{c} -0.5g_1 B_\mu + F_\mu = G_\mu + K_\mu \\ -g_1 B_\mu + F_\mu = G_\mu - K_\mu \end{array} \right|$$

i.e.:

$$\begin{array}{rcl} B_\mu & = & \frac{4}{g_1}K_\mu \\ F_\mu & = & G_\mu + 3K_\mu. \end{array}$$

Let *the charge matrix* be defined as the following:

$$\underline{Y} = -\begin{bmatrix} 1_2 & 0_2 & 0_2 & 0_2 \\ 0_2 & 2\cdot 1_2 & 0_2 & 0_2 \\ 0_2 & 0_2 & 1_2 & 0_2 \\ 0_2 & 0_2 & 0_2 & 2\cdot 1_2 \end{bmatrix}$$

In that case from (9):

$$\begin{array}{c} \underline{\beta_0}i\partial_t\Psi + \underline{\beta_1}i\partial_x\Psi + \underline{\beta_2}i\partial_y\Psi + \underline{\beta_3}i\partial_z\Psi + \\ -h\underline{n}\gamma\Psi - h\underline{k}\beta\Psi + \\ +\left(F_t\underline{\beta_0} + F_x\underline{\beta_1} + F_y\underline{\beta_2} + F_z\underline{\beta_3}\right)\Psi + \\ +0.5g_1\underline{Y}\left(B_t\underline{\beta_0} + B_x\underline{\beta_1} + B_y\underline{\beta_2} + B_z\underline{\beta_3}\right)\Psi = 0. \end{array} \tag{13}$$

Let

$$\begin{gathered}\Psi \rightarrow \Psi‘ = \left(\widetilde{U}\Psi\right),\\ n \rightarrow n‘,\\ k \rightarrow k‘,\\ F_\mu \rightarrow F_\mu‘,\\ B_\mu \rightarrow B_\mu‘\end{gathered}$$

then:

$$\begin{gathered}\underline{\beta_0} i\partial_t \left(\widetilde{U}\Psi\right) + \underline{\beta_1} i\partial_x \left(\widetilde{U}\Psi\right) + \underline{\beta_2} i\partial_y \left(\widetilde{U}\Psi\right) + \underline{\beta_3} i\partial_z \left(\widetilde{U}\Psi\right) +\\ -h\underline{n}‘\gamma\widetilde{U}\Psi - h\underline{k}‘\beta\widetilde{U}\Psi +\\ +\left(F_t‘\underline{\beta_0} + F_x‘\underline{\beta_1} + F_y‘\underline{\beta_2} + F_z‘\underline{\beta_3}\right)\widetilde{U}\Psi +\\ +0.5g_1\underline{Y}\left(B_t‘\underline{\beta_0} + B_x‘\underline{\beta_1} + B_y‘\underline{\beta_2} + B_z‘\underline{\beta_3}\right)\widetilde{U}\Psi = 0\end{gathered}$$

hence:

$$\begin{gathered}\underline{\beta_0} i\left(\partial_t\widetilde{U}\right)\Psi + \underline{\beta_0} i\widetilde{U}\partial_t\Psi + \underline{\beta_1} i\left(\partial_x\widetilde{U}\right)\Psi + \underline{\beta_1} i\widetilde{U}\partial_x\Psi +\\ +\underline{\beta_2} i\left(\partial_y\widetilde{U}\right)\Psi + \underline{\beta_2} i\widetilde{U}\partial_y\Psi + \underline{\beta_3} i\left(\partial_z\widetilde{U}\right)\Psi + \underline{\beta_3} i\widetilde{U}\partial_z\Psi +\\ -h\underline{n}‘\gamma\widetilde{U}\Psi - h\underline{k}‘\beta\widetilde{U}\Psi +\\ +\left(F_t‘\underline{\beta_0} + F_x‘\underline{\beta_1} + F_y‘\underline{\beta_2} + F_z‘\underline{\beta_3}\right)\widetilde{U}\Psi +\\ +0.5g_1\underline{Y}\left(B_t‘\underline{\beta_0} + B_x‘\underline{\beta_1} + B_y‘\underline{\beta_2} + B_z‘\underline{\beta_3}\right)\widetilde{U}\Psi =\\ = 0.\end{gathered}$$

Since

$$\partial_\mu\widetilde{U} = i\frac{\partial_\mu\chi}{2}\begin{bmatrix} \exp\left(i\frac{\chi}{2}\right) & 0 & 0 & 0\\ 0 & 2\exp\left(i\chi\right) & 0 & 0\\ 0 & 0 & \exp\left(i\frac{\chi}{2}\right) & 0\\ 0 & 0 & 0 & 2\exp\left(i\chi\right)\end{bmatrix}$$

then

$$\partial_\mu\widetilde{U} = -i\frac{\partial_\mu\chi}{2}\underline{Y}\widetilde{U};$$

Hence

$$\begin{gathered}
\left(\underline{\beta_0} i\widetilde{U}\partial_t + \underline{\beta_1} i\widetilde{U}\partial_x + \underline{\beta_2} i\widetilde{U}\partial_y + \underline{\beta_3} i\widetilde{U}\partial_z\right)\Psi + \\
-h\left(\underline{n}\text{‘}\gamma\widetilde{U} + \underline{k}\text{‘}\beta\widetilde{U}\right)\Psi + \\
+\left(F_t\text{‘}\underline{\beta_0} + F_x\text{‘}\underline{\beta_1} + F_y\text{‘}\underline{\beta_2} + F_z\text{‘}\underline{\beta_3}\right)\widetilde{U}\Psi + \\
+0.5\left(\begin{array}{c}
\left(g_1\underline{Y}\underline{\beta_0}B_t\text{‘} + \underline{\beta_0}\underline{Y}\partial_t\chi\right) + \left(g_1\underline{Y}\underline{\beta_1}B_x\text{‘} + \underline{\beta_1}\underline{Y}\partial_x\chi\right) + \\
+\left(g_1\underline{Y}\underline{\beta_2}B_y\text{‘} + \underline{\beta_2}\underline{Y}\partial_y\chi\right) + \left(g_1\underline{Y}\underline{\beta_3}B_z\text{‘} + \underline{\beta_3}\underline{Y}\partial_z\chi\right)
\end{array}\right)\widetilde{U}\Psi = \\
= 0
\end{gathered}$$

Since $\underline{Y}\underline{\beta_\mu} = \underline{\beta_\mu}\underline{Y}$ then

$$\begin{gathered}
\left(\underline{\beta_0} i\widetilde{U}\partial_t + \underline{\beta_1} i\widetilde{U}\partial_x + \underline{\beta_2} i\widetilde{U}\partial_y + \underline{\beta_3} i\widetilde{U}\partial_z\right)\Psi + \\
-h\left(\underline{n}\text{‘}\gamma\widetilde{U} + \underline{k}\text{‘}\beta\widetilde{U}\right)\Psi + \\
+\left(F_t\text{‘}\underline{\beta_0} + F_x\text{‘}\underline{\beta_1} + F_y\text{‘}\underline{\beta_2} + F_z\text{‘}\underline{\beta_3}\right)\widetilde{U}\Psi + \\
+0.5\underline{Y}\left(\begin{array}{c}
\underline{\beta_0}\left(g_1B_t\text{‘} + \partial_t\chi\right) + \underline{\beta_1}\left(g_1B_x\text{‘} + \partial_x\chi\right) + \\
+\underline{\beta_2}\left(g_1B_y\text{‘} + \partial_y\chi\right) + \underline{\beta_3}\left(g_1B_z\text{‘} + \partial_z\chi\right)
\end{array}\right)\widetilde{U}\Psi = \\
= 0
\end{gathered}$$

hence:

$$\begin{gathered}
\widetilde{U}^\dagger\left(\underline{\beta_0} i\widetilde{U}\partial_t + \underline{\beta_1} i\widetilde{U}\partial_x + \underline{\beta_2} i\widetilde{U}\partial_y + \underline{\beta_3} i\widetilde{U}\partial_z\right)\Psi + \\
-h\widetilde{U}^\dagger\left(\underline{n}\text{‘}\gamma\widetilde{U} + \underline{k}\text{‘}\beta\widetilde{U}\right)\Psi + \\
+\widetilde{U}^\dagger\left(F_t\text{‘}\underline{\beta_0} + F_x\text{‘}\underline{\beta_1} + F_y\text{‘}\underline{\beta_2} + F_z\text{‘}\underline{\beta_3}\right)\widetilde{U}\Psi + \\
+0.5\widetilde{U}^\dagger\underline{Y}\left(\begin{array}{c}
\underline{\beta_0}\left(g_1B_t\text{‘} + \partial_t\chi\right) + \underline{\beta_1}\left(g_1B_x\text{‘} + \partial_x\chi\right) + \\
+\underline{\beta_2}\left(g_1B_y\text{‘} + \partial_y\chi\right) + \underline{\beta_3}\left(g_1B_z\text{‘} + \partial_z\chi\right)
\end{array}\right)\widetilde{U}\Psi = \\
= 0.
\end{gathered}$$

Because:

$$\begin{gathered}
\widetilde{U}^\dagger\gamma\widetilde{U} = \cos\left(\frac{\chi}{2}\right)\gamma - \sin\left(\frac{\chi}{2}\right)\beta, \\
\widetilde{U}^\dagger\beta\widetilde{U} = \cos\left(\frac{\chi}{2}\right)\beta + \sin\left(\frac{\chi}{2}\right)\gamma, \\
\widetilde{U}^\dagger\widetilde{U} = 1_8, \\
\underline{\beta_\mu}\widetilde{U} = \widetilde{U}\underline{\beta_\mu}, \\
U^\dagger\underline{Y}U = \underline{Y}
\end{gathered}$$

then

$$\begin{gathered}
\left(\underline{\beta_0} i\partial_t + \underline{\beta_1} i\partial_x + \underline{\beta_2} i\partial_y + \underline{\beta_3} i\partial_z\right)\Psi + \\
-h\left(\underline{n}‘\left(\cos\left(\frac{\chi}{2}\right)\gamma - \sin\left(\frac{\chi}{2}\right)\beta\right) + \underline{k}‘\left(\cos\left(\frac{\chi}{2}\right)\beta + \sin\left(\frac{\chi}{2}\right)\gamma\right)\right)\Psi + \\
+\left(F_t‘\underline{\beta_0} + F_x‘\underline{\beta_1} + F_y‘\underline{\beta_2} + F_z‘\underline{\beta_3}\right)\Psi + \\
+0.5\underline{Y}\left(\begin{array}{c} \underline{\beta_0}\left(g_1 B_t‘ + \partial_t\chi\right) + \underline{\beta_1}\left(g_1 B_x‘ + \partial_x\chi\right) + \\ +\underline{\beta_2}\left(g_1 B_y‘ + \partial_y\chi\right) + \underline{\beta_3}\left(g_1 B_z‘ + \partial_z\chi\right) \end{array}\right)\Psi = \\
= 0.
\end{gathered}$$

Therefore from (13):

$$\begin{gathered}
F_x‘ = F_x, \\
B_\mu‘ = B_\mu - \frac{1}{g_1}\partial_\mu\chi, \\
n‘ = -k\sin\frac{\chi}{2} + n\cos\frac{\chi}{2}, \\
k‘ = k\cos\frac{\chi}{2} + n\sin\frac{\chi}{2}.
\end{gathered} \tag{14}$$

But k and n are an integer numbers and $k‘$ and $n‘$ must be an integer numbers, too.

A triplet $\langle l, n, k\rangle$ of integer numbers is *a Fermat triplet* if

$$l^2 = n^2 + k^2.$$

Let ε be any tiny positive real number. An integer number l is *a father number with a precise* ε if for each real number χ and for every Fermat triplet $\langle l, n, k\rangle$ a Fermat triplet $\langle l, n‘, k‘\rangle$ exists for which:

$$\begin{gathered}
\left|-k\sin\frac{\chi}{2} + n\cos\frac{\chi}{2} - n‘\right| < \varepsilon, \\
\left|k\cos\frac{\chi}{2} + n\sin\frac{\chi}{2} - k‘\right| < \varepsilon.
\end{gathered}$$

For every ε*: denumerable many of a father numbers with a precise* ε *exist.*

Excuse me, but I mean that a masses of the real members of the particles families are defined by a father numbers with a precise h. I.e. denumerable many of a families exist.

Therefore for the (14) transformation from (8):

$$\begin{gathered}
\mathbf{\Psi}\left(x_5, x_4\right) = \sum_{r=1}^{4}\phi_r\left(0,0\right)\epsilon_r + \exp\left(-ih\left(nx_5 + kx_4\right)\right)\sum_{k=1}^{4}\phi_k\left(n,k\right)\epsilon_k = \\
= \phi_L\left(0,0\right) + \phi_R\left(0,0\right) + \exp\left(-ih\left(nx_5 + kx_4\right)\right)\left(\phi_L\left(n,k\right) + \phi_R\left(n,k\right)\right) \rightarrow
\end{gathered}$$

$$
\begin{gathered}
\rightarrow \mathbf{\Psi}\text{‘}\left(x_5, x_4\right) = \\
= \exp\left(i\tfrac{\chi}{2}\right)\phi_L\left(0,0\right) + \exp\left(i\chi\right)\phi_R\left(0,0\right) + \\
+ \exp\left(-ih\left(\left(-k\sin\tfrac{\chi}{2} + n\cos\tfrac{\chi}{2}\right)x_5 + \left(k\cos\tfrac{\chi}{2} + n\sin\tfrac{\chi}{2}\right)x_4\right)\right) \cdot \\
\cdot \left(\exp\left(i\tfrac{\chi}{2}\right)\phi_L\left(n,k\right) + \exp\left(i\chi\right)\phi_R\left(n,k\right)\right).
\end{gathered}
$$

$U^{(-)}$ **transformation** $U^{(-)}$ has got the following eigenvalues and eigenvectors:

for the eigenvalue 1: eigenvectors:

$$
\underline{\iota}_1 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \underline{\iota}_2 = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \underline{\iota}_5 = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}, \underline{\iota}_6 = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}; \tag{15}
$$

for the eigenvalue $w = a + i\sqrt{1-a^2}$: eigenvectors:

$$
\underline{\iota}_3 = \frac{1}{\sqrt{2}\sqrt{\sqrt{1-a^2}\left(b+\sqrt{1-a^2}\right)}} \begin{bmatrix} b+\sqrt{1-a^2} \\ 0 \\ 0 \\ 0 \\ ic+g \\ 0 \\ 0 \\ 0 \end{bmatrix}, \tag{16}
$$

$$
\underline{\iota}_4 = \frac{1}{\sqrt{2}\sqrt{\sqrt{1-a^2}\left(b+\sqrt{1-a^2}\right)}} \begin{bmatrix} 0 \\ b+\sqrt{1-a^2} \\ 0 \\ 0 \\ 0 \\ ic+g \\ 0 \\ 0 \end{bmatrix}; \tag{17}
$$

for eigenvalue $w^* = a - i\sqrt{1-a^2}$: eigenvectors:

$$\underline{\iota}_7 = \frac{1}{\sqrt{2}\sqrt{\sqrt{1-a^2}\left(b+\sqrt{1-a^2}\right)}}\begin{bmatrix} ic-g \\ 0 \\ 0 \\ 0 \\ b+\sqrt{1-a^2} \\ 0 \\ 0 \\ 0 \end{bmatrix}, \tag{18}$$

$$\underline{\iota}_8 = \frac{1}{\sqrt{2}\sqrt{\sqrt{1-a^2}\left(b+\sqrt{1-a^2}\right)}}\begin{bmatrix} 0 \\ ic-g \\ 0 \\ 0 \\ 0 \\ b+\sqrt{1-a^2} \\ 0 \\ 0 \end{bmatrix}. \tag{19}$$

Hence the space of $U^{(-)}$ is divided on three orthogonal subspace:
the 4-dimensional $\mathcal{U}_1^{(-)}$on the basis $\langle \underline{\iota}_1, \underline{\iota}_2, \underline{\iota}_5, \underline{\iota}_6 \rangle$ with eigenvalue 1,
the 2-dimensional $\mathcal{U}_w^{(-)}$on the basis $\langle \underline{\iota}_3, \underline{\iota}_4 \rangle$ with eigenvalue w and
the 2-dimensional $\mathcal{U}_{w^*}^{(-)}$on the basis $\langle \underline{\iota}_7, \underline{\iota}_8 \rangle$ with eigenvalue w^*.
Let

$$\widehat{\underline{\iota}}_k = \gamma \underline{\iota}_k.$$

In this case $\langle \widehat{\underline{\iota}}_3, \widehat{\underline{\iota}}_4, \widehat{\underline{\iota}}_7, \widehat{\underline{\iota}}_8 \rangle$ is the orthonormal basis of $\mathcal{U}_1^{(-)}$.

Let $\mathcal{U}_\circ^{(-)}$ be the space on the basis $\langle \widehat{\underline{\iota}}_3, \widehat{\underline{\iota}}_4, \underline{\iota}_3, \underline{\iota}_4 \rangle$ and $\mathcal{U}_*^{(-)}$ be the space on the basis $\langle \widehat{\underline{\iota}}_7, \widehat{\underline{\iota}}_8, \underline{\iota}_7, \underline{\iota}_8 \rangle$.

$$\begin{gathered} \Psi_\circ = \pi_\circ \Psi,\ \Psi_* = \pi_* \Psi, \\ \Psi_\circ \in \mathcal{U}_\circ^{(-)} \text{ and } \Psi_* \in \mathcal{U}_*^{(-)}. \end{gathered} \tag{20}$$

In this case:

$$\pi_\circ = \frac{1}{2\sqrt{1-a^2}}\begin{bmatrix} \left(b+\sqrt{1-a^2}\right)1_4 & (-ic+g)\,\gamma_5 \\ (ic+g)\,\gamma_5 & \left(\sqrt{1-a^2}-b\right)1_4 \end{bmatrix},$$

$$\pi_* = \frac{1}{2\sqrt{1-a^2}} \left[\begin{array}{cc} \left(\sqrt{1-a^2} - b\right) 1_4 & (ic-g)\,\gamma_5 \\ (-g-ic)\,\gamma_5 & \left(b + \sqrt{1-a^2}\right) 1_4 \end{array} \right].$$

Hence

$$\begin{array}{c} \mathbf{\Psi}_\circ\left(x_5, x_4\right) = \frac{1}{2\sqrt{1-a^2}} \cdot \\ \cdot \left(\begin{array}{c} \left(b + \sqrt{(1-a^2)}\right)\left(\phi_L\left(0,0\right) + \phi_R\left(0,0\right)\right) - \\ - \left(ic - g\right)\left(\phi_L\left(n,k\right) - \phi_R\left(n,k\right)\right) + \\ + \exp\left(-ih\left(nx_5 + kx_4\right)\right) \cdot \\ \cdot \left(\begin{array}{c} \left(ic + g\right)\left(\phi_L\left(0,0\right) - \phi_R\left(0,0\right)\right) + \\ + \left(b + \sqrt{(1-a^2)}\right)\left(\phi_L\left(n,k\right) + \phi_R\left(n,k\right)\right) \end{array} \right) \end{array} \right), \end{array} \tag{21}$$

$$\begin{array}{c} \mathbf{\Psi}_*\left(x_5, x_4\right) = \frac{1}{2\sqrt{1-a^2}} \cdot \\ \cdot \left(\begin{array}{c} \left(\sqrt{(1-a^2)} - b\right)\left(\phi_L\left(0,0\right) + \phi_R\left(0,0\right)\right) + \\ + \left(ic - g\right)\left(\phi_L\left(n,k\right) - \phi_R\left(n,k\right)\right) + \\ + \exp\left(-ih\left(nx_5 + kx_4\right)\right) \cdot \\ \cdot \left(\begin{array}{c} \left(g + ic\right)\left(-\phi_L\left(0,0\right) + \phi_R\left(0,0\right)\right) + \\ + \left(b + \sqrt{(1-a^2)}\right)\left(\phi_L\left(n,k\right) + \phi_R\left(n,k\right)\right) \end{array} \right) \end{array} \right). \end{array} \tag{22}$$

If λ is the angle of the $U^{(-)}$ eigenvalue (i.e. $w = a + i\sqrt{1-a^2}$ and $\cos\lambda = a$ and $\sin\lambda = \sqrt{1-a^2}$) then

$$\begin{array}{c} U^{(-)\dagger}\gamma U^{(-)} = \left(\gamma\cos\lambda + \sin\lambda\left(\pi_\circ - \pi_*\right)\beta\right), \\ U^{(-)\dagger}\beta U^{(-)} = \left(\beta\cos\lambda - \sin\lambda\left(\pi_\circ - \pi_*\right)\gamma\right). \end{array} \tag{23}$$

Let

$$\begin{array}{c} \underline{n} \rightarrow \underline{n}\text{‘} = \left(\underline{n}\cos\lambda + \underline{k}\sin\lambda\left(\pi_\circ - \pi_*\right)\right), \\ \underline{k} \rightarrow \underline{k}\text{‘} = \left(\underline{k}\cos\lambda - \underline{n}\sin\lambda\left(\pi_\circ - \pi_*\right)\right), \\ \Psi \rightarrow \Psi\text{‘} = U^{(-)}\Psi, \\ F_\mu \rightarrow F_\mu\text{‘} = F_\mu, \\ B_\mu \rightarrow B_\mu\text{‘} = B_\mu \end{array} \tag{24}$$

and the motion equation for Ψ‘ be (13):

$$\begin{array}{c} \sum_{\mu=0}^{3} \underline{\beta_\mu} i\partial_\mu \Psi\text{‘} - h\left(\underline{n}\text{‘}\gamma + \underline{k}\text{‘}\beta\right)\Psi\text{‘} + \sum_{\mu=0}^{3} F_\mu\text{‘}\underline{\beta_\mu}\Psi\text{‘} + \\ + 0.5 g_1 \underline{Y} \sum_{\mu=0}^{3} B_\mu\text{‘}\underline{\beta_\mu}\Psi\text{‘} = S\Psi\text{‘}. \end{array} \tag{25}$$

Hence:

$$\begin{gathered}
\sum_{\mu=0}^{3}\underline{\beta_\mu}i\left(\partial_\mu U^{(-)}\right)\Psi+\sum_{\mu=0}^{3}\underline{\beta_\mu}iU^{(-)}\left(\partial_\mu\Psi\right)- \\
-h\left(\underline{n}\text{‘}\gamma+\underline{k}\text{‘}\beta\right)\left(U^{(-)}\Psi\right)+\sum_{\mu=0}^{3}F_\mu\text{‘}\underline{\beta_\mu}\left(U^{(-)}\Psi\right)+ \\
+0.5g_1\underline{Y}\sum_{\mu=0}^{3}B_\mu\text{‘}\underline{\beta_\mu}\left(U^{(-)}\Psi\right)=S\left(U^{(-)}\Psi\right),
\end{gathered}$$

then

$$\begin{gathered}
U^{(-)\dagger}\sum_{\mu=0}^{3}\underline{\beta_\mu}i\left(\partial_\mu U^{(-)}\right)\Psi+U^{(-)\dagger}\sum_{\mu=0}^{3}\underline{\beta_\mu}iU^{(-)}\left(\partial_\mu\Psi\right)- \\
-U^{(-)\dagger}h\left(\begin{array}{c}\left(\underline{n}\cos\lambda+\underline{k}\sin\lambda\left(\pi_\circ-\pi_*\right)\right)\gamma+ \\ +\left(\underline{k}\cos\lambda-\underline{n}\sin\lambda\left(\pi_\circ-\pi_*\right)\right)\beta\end{array}\right)U^{(-)}\Psi+ \\
+U^{(-)\dagger}\sum_{\mu=0}^{3}F_\mu\underline{\beta_\mu}U^{(-)}\Psi++U^{(-)\dagger}0.5g_1\underline{Y}\sum_{\mu=0}^{3}B_\mu\underline{\beta_\mu}U^{(-)}\Psi= \\
=U^{(-)\dagger}SU^{(-)}\Psi.
\end{gathered}$$

Since

$$\begin{gathered}
U^{(-)\dagger}\underline{\beta_\mu}=\underline{\beta_\mu}U^{(-)\dagger}, \\
\underline{Y}U^{(-)}=U^{(-)}\underline{Y}, \\
U^{(-)\dagger}\left(\pi_\circ-\pi_*\right)=\left(\pi_\circ-\pi_*\right)U^{(-)\dagger}
\end{gathered}$$

then

$$\begin{gathered}
\sum_{\mu=0}^{3}\underline{\beta_\mu}iU^{(-)\dagger}\left(\partial_\mu U^{(-)}\right)\Psi+\sum_{\mu=0}^{3}\underline{\beta_\mu}i\partial_\mu\Psi- \\
-h\left(\begin{array}{c}\left(\underline{n}\cos\lambda+\underline{k}\sin\lambda\left(\pi_\circ-\pi_*\right)\right)U^{(-)\dagger}\gamma U^{(-)}+ \\ +\left(\underline{k}\cos\lambda-\underline{n}\sin\lambda\left(\pi_\circ-\pi_*\right)\right)U^{(-)\dagger}\beta U^{(-)}\end{array}\right)\Psi+ \\
+\sum_{\mu=0}^{3}F_\mu\underline{\beta_\mu}\Psi+0.5g_1\underline{Y}\sum_{\mu=0}^{3}B_\mu\underline{\beta_\mu}\Psi= \\
=U^{(-)\dagger}SU^{(-)}\Psi.
\end{gathered}$$

From (23):

$$\begin{gathered}
\sum_{\mu=0}^{3}\underline{\beta_\mu}iU^{(-)\dagger}\left(\partial_\mu U^{(-)}\right)\Psi+\sum_{\mu=0}^{3}\underline{\beta_\mu}i\partial_\mu\Psi- \\
-h\left(\begin{array}{c}\left(\underline{n}\cos\lambda+\underline{k}\sin\lambda\left(\pi_\circ-\pi_*\right)\right)\left(\gamma\cos\lambda+\sin\lambda\left(\pi_\circ-\pi_*\right)\beta\right)+ \\ +\left(\underline{k}\cos\lambda-\underline{n}\sin\lambda\left(\pi_\circ-\pi_*\right)\right)\left(\beta\cos\lambda-\sin\lambda\left(\pi_\circ-\pi_*\right)\gamma\right)\end{array}\right)\Psi+ \\
+\sum_{\mu=0}^{3}F_\mu\underline{\beta_\mu}\Psi+0.5g_1\underline{Y}\sum_{\mu=0}^{3}B_\mu\underline{\beta_\mu}\Psi= \\
=U^{(-)\dagger}SU^{(-)}\Psi.
\end{gathered}$$

Since

$$\left(\pi_\circ-\pi_*\right)\left(\pi_\circ-\pi_*\right)=1_8$$

then

$$\begin{gathered}\sum_{\mu=0}^{3}\underline{\beta_{\mu}}iU^{(-)\dagger}\left(\partial_{\mu}U^{(-)}\right)\Psi+\\ +\sum_{\mu=0}^{3}\underline{\beta_{\mu}}i\partial_{\mu}\Psi-h\left(\underline{n}\gamma+\underline{k}\beta\right)\Psi+\sum_{\mu=0}^{3}F_{\mu}\underline{\beta_{\mu}}\Psi+0.5g_1\underline{Y}\sum_{\mu=0}^{3}B_{\mu}\underline{\beta_{\mu}}\Psi=\\ =U^{(-)\dagger}SU^{(-)}\Psi.\end{gathered}$$

Hence from (13):

$$\sum_{\mu=0}^{3}\underline{\beta_{\mu}}iU^{(-)\dagger}\left(\partial_{\mu}U^{(-)}\right)=U^{(-)\dagger}SU^{(-)}$$

and

$$S=\sum_{\mu=0}^{3}\underline{\beta_{\mu}}i\left(\partial_{\mu}U^{(-)}\right)U^{(-)\dagger}$$

Therefore from (25) the motion equation for the transformation (24) is the following:

$$\begin{gathered}\sum_{\mu=0}^{3}\underline{\beta_{\mu}}i\partial_{\mu}\Psi`-\sum_{\mu=0}^{3}\underline{\beta_{\mu}}i\left(\partial_{\mu}U^{(-)}\right)U^{(-)\dagger}\Psi`-\\ -h\left(\underline{n}`\gamma+\underline{k}`\beta\right)\Psi`+\\ +\sum_{\mu=0}^{3}F_{\mu}\underline{\beta_{\mu}}\Psi`+0.5g_1\underline{Y}\sum_{\mu=0}^{3}B_{\mu}\underline{\beta_{\mu}}\Psi`=0.\end{gathered}\tag{26}$$

W**-bosons** Let g_2 be a positive real number.
If design (a,b,c,g form $U^{(-)}$):

$$\begin{aligned}W_{0,\mu}&=-2\frac{1}{g_2g}\left(\begin{array}{c}g\left(\partial_{\mu}a\right)b-g\left(\partial_{\mu}b\right)a+\left(\partial_{\mu}c\right)g^2+\\ +a\left(\partial_{\mu}a\right)c+b\left(\partial_{\mu}b\right)c+c^2\left(\partial_{\mu}c\right)\end{array}\right)\\ W_{1,\mu}&=-2\frac{1}{g_2g}\left(\begin{array}{c}\left(\partial_{\mu}a\right)a^2-bg\left(\partial_{\mu}c\right)+a\left(\partial_{\mu}b\right)b+\\ +a\left(\partial_{\mu}c\right)c+g^2\left(\partial_{\mu}a\right)+c\left(\partial_{\mu}b\right)g\end{array}\right)\\ W_{2,\mu}&=-2\frac{1}{g_2g}\left(\begin{array}{c}g\left(\partial_{\mu}a\right)c-a\left(\partial_{\mu}a\right)b-b^2\left(\partial_{\mu}b\right)-\\ -c\left(\partial_{\mu}c\right)b-\left(\partial_{\mu}b\right)g^2-\left(\partial_{\mu}c\right)ga\end{array}\right)\end{aligned}$$

and

$$W_{\mu}=\left[\begin{array}{cccc}W_{0,\mu}1_2&0_2&\left(W_{1,\mu}-iW_{2,\mu}\right)1_2&0_2\\ 0_2&0_2&0_2&0_2\\ \left(W_{1,\mu}+iW_{2,\mu}\right)1_2&0_2&-W_{0,\mu}1_2&0_2\\ 0_2&0_2&0_2&0_2\end{array}\right]$$

then

$$-\, i\left(\partial_\mu U^{(-)}\right) U^{(-)\dagger}=\frac{1}{2}g_2 W_\mu \tag{27}$$

and from (26):

$$\begin{array}{c}\sum_{\mu=0}^{3}\underline{\beta_\mu} i\left(\partial_\mu - i\frac{1}{2}g_2 W_\mu\right)\Psi\text{‘}- \\ -h\left(\underline{n}\text{‘}\gamma+\underline{k}\text{‘}\beta\right)\Psi\text{‘}+ \\ +\sum_{\mu=0}^{3}F_\mu\underline{\beta_\mu}\Psi\text{‘}+0.5g_1\underline{Y}\sum_{\mu=0}^{3}B_\mu\underline{\beta_\mu}\Psi\text{‘}=0.\end{array} \tag{28}$$

Let

$$\overset{,}{U}=\left[\begin{array}{cccc}\left(\overset{,}{a}+i\overset{,}{b}\right)1_2 & 0_2 & \left(\overset{,}{c}+i\overset{,}{g}\right)1_2 & 0_2 \\ 0_2 & 1_2 & 0_2 & 0_2 \\ \left(-\overset{,}{c}+i\overset{,}{g}\right)1_2 & 0_2 & \left(\overset{,}{a}-i\overset{,}{b}\right)1_2 & 0_2 \\ 0_2 & 0_2 & 0_2 & 1_2\end{array}\right],$$

$$\overset{,}{\pi}_\circ=\frac{1}{2\sqrt{1-\overset{,}{a}^2}}\left[\begin{array}{cc}\left(\overset{,}{b}+\sqrt{1-\overset{,}{a}^2}\right)1_4 & \left(-i\overset{,}{c}+\overset{,}{g}\right)\gamma_5 \\ \left(i\overset{,}{c}+\overset{,}{g}\right)\gamma_5 & \left(\sqrt{1-\overset{,}{a}^2}-\overset{,}{b}\right)1_4\end{array}\right],$$

$$\overset{,}{\pi}_*=\frac{1}{2\sqrt{1-\overset{,}{a}^2}}\left[\begin{array}{cc}\left(\sqrt{1-\overset{,}{a}^2}-\overset{,}{b}\right)1_4 & \left(i\overset{,}{c}-\overset{,}{g}\right)\gamma_5 \\ \left(-\overset{,}{g}-i\overset{,}{c}\right)\gamma_5 & \left(\overset{,}{b}+\sqrt{1-\overset{,}{a}^2}\right)1_4\end{array}\right].$$

Let:

$$\cos\overset{,}{\lambda}=\overset{,}{a}\ \text{ and }\ \sin\overset{,}{\lambda}=\sqrt{1-\overset{,}{a}^2}$$

and

$$\begin{array}{c}\Psi\text{‘}\rightarrow\Psi'=\left(\overset{,}{U}\,\Psi\text{‘}\right), \\ \underline{n}\text{‘}\rightarrow\underline{n}'=\left(\underline{n}\text{‘}\cos\overset{,}{\lambda}+\underline{k}\text{‘}\sin\overset{,}{\lambda}\left(\overset{,}{\pi}_\circ-\overset{,}{\pi}_*\right)\right), \\ \underline{k}\text{‘}\rightarrow\underline{k}'=\left(\underline{k}\text{‘}\cos\overset{,}{\lambda}-\underline{n}\text{‘}\sin\overset{,}{\lambda}\left(\overset{,}{\pi}_\circ-\overset{,}{\pi}_*\right)\right), \\ F_\mu\rightarrow F'_\mu=F_\mu, \\ B_\mu\rightarrow B'_\mu=B_\mu,\end{array} \tag{29}$$

and

$$W_\mu \to W'_\mu.$$

In that case from (27):

$$W'_\mu = -\frac{2i}{g_2}\left(\partial_\mu\left(\acute{U}\,U^{(-)}\right)\right)\left(\acute{U}\,U^{(-)}\right)^\dagger;$$

Hence:

$$W'_\mu = -\frac{2i}{g_2}\left(\partial_\mu\,\acute{U}\right)\acute{U}^\dagger - \frac{2i}{g_2}\,\acute{U}\,\left(\partial_\mu U^{(-)}\right)U^{(-)\dagger}\,\acute{U}^\dagger\,;$$

i.e.:

$$W'_\mu = \acute{U}\,W_\mu\,\acute{U}^\dagger - \frac{2i}{g_2}\left(\partial_\mu\,\acute{U}\right)\acute{U}^\dagger\,.$$

If

$$F_{\mu,\nu} = \left(\partial_\mu W_\nu - \partial_\nu W_\mu - i\frac{g_2}{2}\left(W_\mu W_\nu - W_\nu W_\mu\right)\right)$$

then

$$F'_{\mu,\nu} = \partial_\mu W'_\nu - \partial_\nu W'_\mu - i\frac{g_2}{2}\left(W'_\mu W'_\nu - W'_\nu W'_\mu\right) = UF_{\mu,\nu}U^\dagger.$$

Therefore $F_{\mu,\nu}$ is invariant for the transformation (29).
The Lagrangian for $F_{\mu,\nu}$:

$$\mathcal{L}_F = \left(-\frac{1}{4}\sum_{\mu,\nu} F^{\mu,\nu}F_{\mu,\nu}\right).$$

Hence the Euler-Lagrange equations for W_μ are the following:

$$\sum_\nu \partial^\nu\left(\partial_\mu W_\nu - \partial_\nu W_\mu - i\frac{g_2}{2}\left[W_\mu, W_\nu\right]\right) = 0.$$

For the components:

$$\begin{array}{l}
\sum_\nu \partial^\nu\partial_\nu W_{0,\mu} = g_2\sum_\nu \partial^\nu\left(W_{1,\mu}W_{2,\nu} - W_{2,\mu}W_{1,\nu}\right) + \partial_\mu\sum_\nu \partial^\nu W_{0,\nu},\\
\sum_\nu \partial^\nu\partial_\nu W_{1,\mu} = g_2\sum_\nu \partial^\nu\left(W_{0,\nu}W_{2,\mu} - W_{0,\mu}W_{2,\nu}\right) + \partial_\mu\sum_\nu \partial^\nu W_{1,\nu},\\
\sum_\nu \partial^\nu\partial_\nu W_{2,\mu} = g_2\sum_\nu \partial^\nu\left(W_{0,\mu}W_{1,\nu} - W_{0,\nu}W_{1,\mu}\right) + \partial_\mu\sum_\nu \partial^\nu W_{2,\nu}.
\end{array}$$

Let:

$$\begin{array}{c}\alpha_{0,\mu,\nu} = \partial_\nu W_{0,\mu} - g_2 \left(W_{1,\mu} W_{2,\nu} - W_{2,\mu} W_{1,\nu}\right), \\ \alpha_{1,\mu,\nu} = \partial_\nu W_{1,\mu} - g_2 \left(W_{0,\nu} W_{2,\mu} - W_{0,\mu} W_{2,\nu}\right), \\ \alpha_{2,\mu,\nu} = \partial_\nu W_{2,\mu} - g_2 \left(W_{0,\mu} W_{1,\nu} - W_{0,\nu} W_{1,\mu}\right).\end{array} \tag{30}$$

Hence if $\sum_\nu \partial^\nu W_\nu = 0$ then

$$\sum_\nu \partial^\nu \alpha_{0,\mu,\nu} = 0, \sum_\nu \partial^\nu \alpha_{1,\mu,\nu} = 0, \sum_\nu \partial^\nu \alpha_{2,\mu,\nu} = 0.$$

From (30):

$$\partial_\nu W_{0,\mu} = \left(g_2 \left(W_{1,\mu} W_{2,\nu} - W_{2,\mu} W_{1,\nu}\right) + \alpha_{0,\mu,\nu}\right), \tag{31}$$

$$\partial_\nu W_{1,\mu} = \left(g_2 \left(W_{0,\nu} W_{2,\mu} - W_{0,\mu} W_{2,\nu}\right) + \alpha_{1,\mu,\nu}\right), \tag{32}$$

$$\partial_\nu W_{2,\mu} = \left(g_2 \left(W_{0,\mu} W_{1,\nu} - W_{0,\nu} W_{1,\mu}\right) + \alpha_{2,\mu,\nu}\right); \tag{33}$$

from (31):

$$\begin{array}{c}\partial_\nu \partial_\nu W_{0,\mu} = g_2 \partial_\nu \left(W_{1,\mu} W_{2,\nu} - W_{2,\mu} W_{1,\nu}\right) + \partial_\nu \alpha_{0,\mu,\nu} = \\ = g_2 \left(\partial_\nu W_{1,\mu} W_{2,\nu} + W_{1,\mu} \partial_\nu W_{2,\nu} - \partial_\nu W_{2,\mu} W_{1,\nu} - W_{2,\mu} \partial_\nu W_{1,\nu}\right) + \partial_\nu \alpha_{0,\mu,\nu};\end{array} \tag{34}$$

hence from (34), (32) and (33):

$$\begin{array}{rcl}\partial_\nu \partial_\nu W_{0,\mu} & = & g_2 \left(\begin{array}{c}\left(g_2 \left(W_{0,\nu} W_{2,\mu} - W_{0,\mu} W_{2,\nu}\right) + \alpha_{1,\mu,\nu}\right) W_{2,\nu} - \\ - \left(g_2 \left(W_{0,\mu} W_{1,\nu} - W_{0,\nu} W_{1,\mu}\right) + \alpha_{2,\mu,\nu}\right) W_{1,\nu} - \\ - W_{2,\mu} \partial_\nu W_{1,\nu} + W_{1,\mu} \partial_\nu W_{2,\nu}\end{array}\right) + \\ & & + \partial_\nu \alpha_{0,\mu,\nu};\end{array}$$

hence:

$$\begin{array}{c}\partial_\nu \partial_\nu W_{0,\mu} = \\ = g_2 \left(\begin{array}{c}g_2 \left(-\left(W_{2,\nu}^2 + W_{1,\nu}^2\right) W_{0,\mu} + \left(W_{1,\mu} W_{1,\nu} + W_{2,\mu} W_{2,\nu}\right) W_{0,\nu}\right) + \\ + \alpha_{1,\mu,\nu} W_{2,\nu} - \alpha_{2,\mu,\nu} W_{1,\nu} + W_{1,\mu} \partial_\nu W_{2,\nu} - W_{2,\mu} \partial_\nu W_{1,\nu}\end{array}\right) + \\ + \partial_\nu \alpha_{0,\mu,\nu};\end{array}$$

and

$$\begin{gathered}\partial_\nu \partial_\nu W_{0,\mu} = -g_2^2 \left(W_{2,\nu}^2 + W_{1,\nu}^2 + W_{0,\nu}^2\right) W_{0,\mu} + \\ + g_2^2 \left(W_{0,\mu} W_{0,\nu} + W_{1,\mu} W_{1,\nu} + W_{2,\mu} W_{2,\nu}\right) W_{0,\nu} + \\ + g_2 \left(\alpha_{1,\mu,\nu} W_{2,\nu} - \alpha_{2,\mu,\nu} W_{1,\nu} + W_{1,\mu} \partial_\nu W_{2,\nu} - W_{2,\mu} \partial_\nu W_{1,\nu}\right) + \partial_\nu \alpha_{0,\mu,\nu};\end{gathered}$$

if $\sum_\nu \partial_\nu W_\nu = 0$ then:

$$\begin{gathered}\sum_\nu \partial_\nu \partial_\nu W_{0,\mu} = -g_2^2 W_{0,\mu} \sum_\nu W_\nu^2 + \\ + \frac{g_2^2}{2} \sum_\nu \left(W_\mu W_\nu + W_\nu W_\mu\right) W_{0,\nu} + g_2 \sum_\nu \left(\alpha_{1,\mu,\nu} W_{2,\nu} - \alpha_{2,\mu,\nu} W_{1,\nu}\right),\end{gathered} \tag{35}$$

$$\begin{gathered}\sum_\nu \partial_\nu \partial_\nu W_{1,\mu} = -g_2^2 W_{1,\mu} \sum_\nu W_\nu^2 + \\ + \frac{g_2^2}{2} \sum_\nu \left(W_\nu W_\mu + W_\mu W_\nu\right) W_{1,\nu} + g_2 \sum_\nu \left(W_{0,\nu} \alpha_{2,\mu,\nu} - \alpha_{0,\mu,\nu} W_{2,\nu}\right)\end{gathered}$$

and

$$\begin{gathered}\sum_\nu \partial_\nu \partial_\nu W_{2,\mu} = -g_2^2 W_{2,\mu} \sum_\nu W_\nu^2 + \\ + \frac{g_2^2}{2} \sum_\nu \left(W_\nu W_\mu + W_\mu W_\nu\right) W_{2,\nu} + g_2 \sum_\nu \left(\alpha_{0,\mu,\nu} W_{1,\nu} - W_{0,\nu} \alpha_{1,\mu,\nu}\right).\end{gathered}$$

$$\alpha_{\mu,\nu} = \left[\begin{array}{cc} \alpha_{0,\mu,\nu} & \alpha_{1,\mu,\nu} - i\alpha_{2,\mu,\nu} \\ \alpha_{1,\mu,\nu} + i\alpha_{2,\mu,\nu} & -\alpha_{0,\mu,\nu} \end{array}\right]$$

then

$$\begin{gathered}\sum_\nu \partial_\nu \partial_\nu W_\mu = -g_2^2 W_\mu \sum_\nu W_\nu^2 + \\ + \frac{g_2^2}{2} \sum_\nu \left(W_\nu W_\mu + W_\mu W_\nu\right) W_\nu - i\frac{g_2^2}{2} \sum_\nu \left(\alpha_{\mu,\nu} W_\nu - W_\nu \alpha_{\mu,\nu}\right).\end{gathered}$$

It is the motion equation for the field W_μ which has got a less than unit 1 velocity. That is this field does not behave as a massless field.

Hence although $F_{\mu,\nu}$ is a massless field but its components W_μ do not behave like a massless fields.

If

$$\sum_\nu \left(W_\nu \frac{\partial W_\nu}{\partial W_\mu} + \frac{\partial W_\nu}{\partial W_\mu} W_\nu\right) = 0$$

then a real υ exists for which

$$\upsilon = \left(2\sum_{\nu} W_{\nu}^{2} \right)^{\frac{1}{2}} \tag{36}$$

and

$$\partial_{W_{\mu}} \upsilon = 0$$

then the Lagrangian of W_{μ} is:

$$\begin{gathered}\widehat{\mathcal{L}} = \textstyle\sum_{\nu} \left(\partial_{\nu} W_{\mu}\right) \left(\partial_{\nu} W_{\mu}\right) - g_2^2 \frac{\upsilon^2}{2} W_{\mu}^2 + \\ + \frac{g_2^2}{4} \textstyle\sum_{\nu} \left(W_{\nu} W_{\mu} + W_{\mu} W_{\nu}\right)^2 - \\ - i \frac{g_2^2}{2} \left(\left(\textstyle\sum_{\nu} \left[\alpha_{\mu,\nu}, W_{\nu}\right] \right) W_{\mu} + W_{\mu} \left(\textstyle\sum_{\nu} \left[\alpha_{\mu,\nu}, W_{\nu}\right] \right) \right).\end{gathered}$$

It is a lagrangian of a field with mass

$$M = g_2 \frac{\upsilon}{\sqrt{2}}$$

and $M > 0$.

A **and** *Z* **bosons** Let A_{μ} and Z_{μ} are a fields for which [7]:

$$Z_{\mu} = \frac{1}{\sqrt{g_1^2 + g_2^2}} \left(g_2 W_{0,\mu} - g_1 B_{\mu}\right), \, A_{\mu} = \frac{1}{\sqrt{g_1^2 + g_2^2}} \left(g_2 B_{\mu} + g_1 W_{0,\mu}\right) \tag{37}$$

and

$$\sum_{\nu} \partial^{\nu} \partial_{\nu} A_{\mu} = 0. \tag{38}$$

Let denote:

$$\frac{g_2^2}{2} \sum_{\nu} \left(W_{\mu} W_{\nu} + W_{\nu} W_{\mu}\right) W_{0,\nu} + g_2 \sum_{\nu} \left(\alpha_{1,\mu,\nu} W_{2,\nu} - \alpha_{2,\mu,\nu} W_{1,\nu}\right) = \Lambda.$$

Hence from (35) and (36):

$$\sum_{\nu} \partial_{\nu} \partial_{\nu} W_{0,\mu} = -g_2^2 \frac{\upsilon^2}{2} W_{0,\mu} + \Lambda \tag{39}$$

From (37):

$$B_\mu = \frac{1}{\sqrt{g_1^2+g_2^2}}\left(g_2 A_\mu - g_1 Z_\mu\right),\ W^0_\mu = \frac{1}{\sqrt{g_1^2+g_2^2}}\left(g_1 A_\mu + g_2 Z_\mu\right). \tag{40}$$

and

$$\sum_\nu \partial^\nu \partial_\nu A_\mu = \frac{1}{\sqrt{g_1^2+g_2^2}}\left(g_2 \sum_\nu \partial^\nu \partial_\nu B_\mu + g_1 \sum_\nu \partial^\nu \partial_\nu W_{0,\mu}\right),$$

from (39):

$$\sum_\nu \partial^\nu \partial_\nu A_\mu = \frac{1}{\sqrt{g_1^2+g_2^2}}\left(\begin{array}{c} g_2\left(\sum_\nu \partial^\nu \partial_\nu B_\mu + g_1^2 \frac{\upsilon^2}{2} B_\mu - g_1^2 \frac{\upsilon^2}{2} B_\mu\right) + \\ + g_1\left(-g_2^2 \frac{\upsilon^2}{2} W_{0,\mu} + \Lambda\right)\end{array}\right),$$

from (40)

$$\begin{aligned}\sum_\nu \partial^\nu \partial_\nu A_\mu \quad &= \quad -\frac{\upsilon^2}{2} g_1 g_2 \frac{1}{g_1^2+g_2^2}\left(2 g_1 g_2 A_\mu + \left(g_2^2 - g_1^2\right) Z_\mu\right) + \\ &\quad + \frac{1}{\sqrt{g_1^2+g_2^2}}\left(g_1 \Lambda + g_2\left(\sum_\nu \partial^\nu \partial_\nu B_\mu + g_1^2 \frac{\upsilon^2}{2} B_\mu\right)\right),\end{aligned}$$

from (38):

$$\begin{array}{c} A_\mu = -\left(g_2^2 - g_1^2\right) \frac{1}{2 g_1 g_2} Z_\mu + \\ + \frac{1}{\upsilon^2 (g_1 g_2)^2} \sqrt{g_1^2+g_2^2}\left(g_1 \Lambda + g_2\left(\sum_\nu \partial^\nu \partial_\nu B_\mu + g_1^2 \frac{\upsilon^2}{2} B_\mu\right)\right)\end{array} \tag{41}$$

From (37):

$$\sum_\nu \partial^\nu \partial_\nu Z_\mu = \frac{1}{\sqrt{g_1^2+g_2^2}}\left(g_2 \sum_\nu \partial^\nu \partial_\nu W_{0,\mu} - g_1 \sum_\nu \partial^\nu \partial_\nu B_\mu\right),$$

from (39):

$$\sum_\nu \partial^\nu \partial_\nu Z_\mu = \frac{1}{\sqrt{g_1^2+g_2^2}}\left(\begin{array}{c} g_2\left(-g_2^2 \frac{\upsilon^2}{2} W_{0,\mu} + \Lambda\right) - \\ - g_1\left(\sum_\nu \partial^\nu \partial_\nu B_\mu + g_1^2 \frac{\upsilon^2}{2} B_\mu - g_1^2 \frac{\upsilon^2}{2} B_\mu\right)\end{array}\right),$$

from (40):

$$\sum_{\nu}\partial^{\nu}\partial_{\nu}Z_{\mu}=-\frac{\upsilon^{2}}{2}\frac{1}{g_{1}^{2}+g_{2}^{2}}\left(g_{1}^{4}+g_{2}^{4}\right)Z_{\mu}-g_{1}g_{2}\frac{\upsilon^{2}}{2}\frac{1}{g_{1}^{2}+g_{2}^{2}}\left(g_{2}^{2}-g_{1}^{2}\right)A_{\mu}+$$

$$+\frac{1}{\sqrt{g_{1}^{2}+g_{2}^{2}}}\left(g_{2}\Lambda-g_{1}\left(\sum_{\nu}\partial^{\nu}\partial_{\nu}B_{\mu}+g_{1}^{2}\frac{\upsilon^{2}}{2}B_{\mu}\right)\right)$$

and from (41):

$$\begin{array}{c}\sum_{\nu}\partial^{\nu}\partial_{\nu}Z_{\mu}=-\frac{1}{2}\frac{\upsilon^{2}}{2}\left(g_{1}^{2}+g_{2}^{2}\right)Z_{\mu}+\\ +\frac{1}{2}\sqrt{g_{1}^{2}+g_{2}^{2}}\left(\frac{1}{g_{2}}\Lambda-\frac{1}{g_{1}}\left(\sum_{\nu}\partial^{\nu}\partial_{\nu}B_{\mu}+g_{1}^{2}\frac{\upsilon^{2}}{2}B_{\mu}\right)\right)\end{array}$$

That is Z_{μ} has got the mass:

$$M_{Z}=\frac{\upsilon}{2}\sqrt{g_{1}^{2}+g_{2}^{2}}.$$

Fragments Since

$$\left(\pi_{\circ}+\pi_{*}\right)=1_{8}$$

then

$$\begin{array}{c}\sum_{\mu=0}^{3}\underline{\beta_{\mu}}i\left(\partial_{\mu}-i\frac{1}{2}g_{2}W_{\mu}\right)\left(\pi_{\circ}+\pi_{*}\right)\Psi`-\\ -h\left(\begin{array}{c}\left(\underline{n}\cos\lambda\left(\pi_{\circ}+\pi_{*}\right)+\underline{k}\sin\lambda\left(\pi_{\circ}-\pi_{*}\right)\right)\gamma+\\ +\left(\underline{k}\cos\lambda\left(\pi_{\circ}+\pi_{*}\right)-\underline{n}\sin\lambda\left(\pi_{\circ}-\pi_{*}\right)\right)\beta\end{array}\right)\Psi`+\\ +\sum_{\mu=0}^{3}F_{\mu}\underline{\beta_{\mu}}\left(\pi_{\circ}+\pi_{*}\right)\Psi`+\\ +0.5g_{1}\underline{Y}\sum_{\mu=0}^{3}\overline{B_{\mu}}\underline{\beta_{\mu}}\left(\pi_{\circ}+\pi_{*}\right)\Psi`=0.\end{array}$$

Because

$$\begin{array}{c}\pi_{\circ}\beta=\beta\pi_{\circ},\ \pi_{*}\beta=\beta\pi_{*},\\ \pi_{\circ}\gamma=\gamma\pi_{\circ},\pi_{*}\gamma=\gamma\pi_{*}\end{array}$$

then

$$\begin{gathered}
\sum_{\mu=0}^{3}\underline{\beta_\mu}i\left(\partial_\mu - i\frac{1}{2}g_2W_\mu\right)\pi_\circ\Psi`+\\
+\sum_{\mu=0}^{3}\underline{\beta_\mu}i\left(\partial_\mu - i\frac{1}{2}g_2W_\mu\right)\pi_*\Psi`-\\
-h\left(\begin{array}{c}
\underline{n}\cos\lambda\gamma\pi_\circ\Psi` + \underline{n}\cos\lambda\gamma\pi_*\Psi`+\\
+\underline{k}\sin\lambda\gamma\pi_\circ\Psi` - \underline{k}\sin\lambda\gamma\pi_*\Psi`+\\
+\underline{k}\cos\lambda\beta\pi_\circ\Psi` + \underline{k}\cos\lambda\beta\pi_*\Psi`-\\
-\underline{n}\sin\lambda\beta\pi_\circ\Psi` + \underline{n}\sin\lambda\beta\pi_*\Psi`
\end{array}\right)+\\
+\sum_{\mu=0}^{3}F_\mu\underline{\beta_\mu}\pi_\circ\Psi` + \sum_{\mu=0}^{3}F_\mu\underline{\beta_\mu}\pi_*\Psi`+\\
+0.5g_1\underline{Y}\sum_{\mu=0}^{3}B_\mu\underline{\beta_\mu}\pi_\circ\Psi`+\\
+0.5g_1\underline{Y}\sum_{\mu=0}^{3}B_\mu\underline{\beta_\mu}\pi_*\Psi` = 0.
\end{gathered}$$

Let

$$\Psi_\circ` = \pi_\circ\Psi` \text{ and } \Psi_*` = \pi_*\Psi`.$$

In that case:

$$\begin{gathered}
\sum_{\mu=0}^{3}\underline{\beta_\mu}i\left(\partial_\mu - i\frac{1}{2}g_2W_\mu\right)\Psi_\circ`-\\
-h\left(\left(\underline{n}\cos\lambda + \underline{k}\sin\lambda\right)\gamma + \left(\underline{k}\cos\lambda - \underline{n}\sin\lambda\right)\beta\right)\Psi_\circ`+\\
+\sum_{\mu=0}^{3}F_\mu\underline{\beta_\mu}\Psi_\circ` + 0.5g_1\underline{Y}\sum_{\mu=0}^{3}B_\mu\underline{\beta_\mu}\Psi_\circ`+\\
\sum_{\mu=0}^{3}\underline{\beta_\mu}i\left(\partial_\mu - i\frac{1}{2}g_2W_\mu\right)\Psi_*`-\\
-h\left(\left(\underline{n}\cos\lambda - \underline{k}\sin\lambda\right)\gamma + \left(\underline{k}\cos\lambda + \underline{n}\sin\lambda\right)\beta\right)\Psi_*`+\\
+\sum_{\mu=0}^{3}F_\mu\underline{\beta_\mu}\Psi_*` + 0.5g_1\underline{Y}\sum_{\mu=0}^{3}B_\mu\underline{\beta_\mu}\Psi_*` = 0.
\end{gathered} \tag{42}$$

Therefore for the (24) transformation from (21, 22):

$$\mathbf{\Psi}\left(x_5, x_4\right) = \mathbf{\Psi}_\circ\left(x_5, x_4\right) + \mathbf{\Psi}_*\left(x_5, x_4\right) \rightarrow$$

$$\rightarrow \mathbf{\Psi}`\left(x_5, x_4\right) = \mathbf{\Psi}_\circ`\left(x_5, x_4\right) + \mathbf{\Psi}_*`\left(x_5, x_4\right) =$$

$$= \frac{1}{2\sqrt{1-a^2}}\cdot$$

$$\cdot\left(\begin{array}{c}
\left(b + \sqrt{(1-a^2)}\right)\left(\left(a + i\sqrt{1-a^2}\right)\phi_L(0,0) + \phi_R(0,0)\right) -\\
-\left(ic - g\right)\left(\left(a + i\sqrt{1-a^2}\right)\phi_L(n,k) - \phi_R(n,k)\right) +\\
+\exp\left(-ih\left(\left(n\cos\lambda + k\sin\lambda\right)x_5 + \left(k\cos\lambda - n\sin\lambda\right)x_4\right)\right)\cdot\\
\cdot\left(\begin{array}{c}
\left(ic + g\right)\left(\left(a + i\sqrt{1-a^2}\right)\phi_L(0,0) - \phi_R(0,0)\right) +\\
+\left(b + \sqrt{(1-a^2)}\right)\left(\left(a + i\sqrt{1-a^2}\right)\phi_L(n,k) + \phi_R(n,k)\right)
\end{array}\right)
\end{array}\right)+$$

$$+\frac{1}{2\sqrt{1-a^2}}\cdot$$

$$\cdot\left(\begin{array}{c}\left(\sqrt{(1-a^2)}-b\right)\left(\left(a-i\sqrt{1-a^2}\right)\phi_L(0,0)+\phi_R(0,0)\right)+\\ +(ic-g)\left(\left(a-i\sqrt{1-a^2}\right)\phi_L(n,k)-\phi_R(n,k)\right)+\\ +\exp\left(-ih\left((n\cos\lambda-k\sin\lambda)x_5+(k\cos\lambda+n\sin\lambda)x_4\right)\right)\cdot\\ \cdot\left(\begin{array}{c}(g+ic)\left(-\left(a-i\sqrt{1-a^2}\right)\phi_L(0,0)+\phi_R(0,0)\right)+\\ +\left(b+\sqrt{(1-a^2)}\right)\left(\left(a-i\sqrt{1-a^2}\right)\phi_L(n,k)+\phi_R(n,k)\right)\end{array}\right)\end{array}\right).$$

That is:

$$\boldsymbol{\Psi}(x_5,x_4)=\boldsymbol{\Psi}_{\circ}(x_5,x_4)+\boldsymbol{\Psi}_{*}(x_5,x_4)\rightarrow$$

$$\rightarrow\boldsymbol{\Psi}\text{‘}(x_5,x_4)=\boldsymbol{\Psi}_{\circ}\text{‘}(x_5,x_4)+\boldsymbol{\Psi}_{*}\text{‘}(x_5,x_4)=$$

$$=\frac{1}{2\sqrt{1-a^2}}\cdot$$

$$\cdot\left(\begin{array}{c}\left(b+\sqrt{(1-a^2)}\right)\left(\left(a+i\sqrt{1-a^2}\right)\phi_L(0,0)+\phi_R(0,0)\right)-\\ -(ic-g)\left(\left(a+i\sqrt{1-a^2}\right)\phi_L(n,k)-\phi_R(n,k)\right)+\\ +\exp\left(-ih\left(\left(na+k\sqrt{1-a^2}\right)x_5+\left(ka-n\sqrt{1-a^2}\right)x_4\right)\right)\cdot\\ \cdot\left(\begin{array}{c}(ic+g)\left(\left(a+i\sqrt{1-a^2}\right)\phi_L(0,0)-\phi_R(0,0)\right)+\\ +\left(b+\sqrt{(1-a^2)}\right)\left(\left(a+i\sqrt{1-a^2}\right)\varphi_L(n,k)+\phi_R(n,k)\right)\end{array}\right)\end{array}\right)+$$

$$+\frac{1}{2\sqrt{1-a^2}}\cdot$$

$$\cdot\left(\begin{array}{c}\left(\sqrt{(1-a^2)}-b\right)\left(\left(a-i\sqrt{1-a^2}\right)\varphi_L(0,0)+\phi_R(0,0)\right)+\\ +(ic-g)\left(\left(a-i\sqrt{1-a^2}\right)\phi_L(n,k)-\phi_R(n,k)\right)+\\ +\exp\left(-ih\left(\left(na-k\sqrt{1-a^2}\right)x_5+\left(ka+n\sqrt{1-a^2}\right)x_4\right)\right)\cdot\\ \cdot\left(\begin{array}{c}(g+ic)\left(-\left(a-i\sqrt{1-a^2}\right)\phi_L(0,0)+\phi_R(0,0)\right)+\\ +\left(b+\sqrt{(1-a^2)}\right)\left(\left(a-i\sqrt{1-a^2}\right)\phi_L(n,k)+\phi_R(n,k)\right)\end{array}\right)\end{array}\right).$$

Let in some point $\langle t,x,y,z\rangle$ $\phi_L(n,k)\neq\mathbf{0}$ or/and $\phi_R(n,k)\neq\mathbf{0}$
In that case (7) in this point: $\phi_L(0,0)=\mathbf{0}$ and $\phi_R(0,0)=\mathbf{0}$.
Hence:

$$\begin{gathered}\mathbf{\Psi}_{\circ}'(x_5,x_4)=\frac{1}{2\sqrt{1-a^2}}\cdot\\ \cdot\left(\begin{array}{c}-(ic-g)\left(\left(a+i\sqrt{1-a^2}\right)\phi_L(n,k)-\phi_R(n,k)\right)+\\ +\exp\left(-ih\left(\left(na+k\sqrt{1-a^2}\right)x_5+\left(ka-n\sqrt{1-a^2}\right)x_4\right)\right)\cdot\\ \cdot\left(b+\sqrt{(1-a^2)}\right)\left(\left(a+i\sqrt{1-a^2}\right)\phi_L(n,k)+\phi_R(n,k)\right)\end{array}\right),\end{gathered}\tag{43}$$

$$\begin{gathered}\mathbf{\Psi}_{*}'(x_5,x_4)=\frac{1}{2\sqrt{1-a^2}}\cdot\\ \cdot\left(\begin{array}{c}(ic-g)\left(\left(a-i\sqrt{1-a^2}\right)\phi_L(n,k)-\phi_R(n,k)\right)+\\ +\exp\left(-ih\left(\left(na-k\sqrt{1-a^2}\right)x_5+\left(ka+n\sqrt{1-a^2}\right)x_4\right)\right)\cdot\\ \cdot\left(b+\sqrt{(1-a^2)}\right)\left(\left(a-i\sqrt{1-a^2}\right)\phi_L(n,k)+\phi_R(n,k)\right)\end{array}\right)\end{gathered}\tag{44}$$

and

$$\begin{gathered}\mathbf{\Psi}'(x_5,x_4)=\\ =-i(ic-g)\phi_L(n,k)+\\ +\frac{1}{2\sqrt{1-a^2}}\cdot\\ \cdot(\exp\left(-ih\left(\left(na+k\sqrt{1-a^2}\right)x_5+\left(ka-n\sqrt{1-a^2}\right)x_4\right)\right)\cdot\\ \cdot\left(b+\sqrt{(1-a^2)}\right)\left(\left(a+i\sqrt{1-a^2}\right)\phi_L(n,k)+\phi_R(n,k)\right)+\\ +\frac{1}{2\sqrt{1-a^2}}\cdot\\ +\exp\left(-ih\left(\left(na-k\sqrt{1-a^2}\right)x_5+\left(ka+n\sqrt{1-a^2}\right)x_4\right)\right)\cdot\\ \cdot\left(b+\sqrt{(1-a^2)}\right)\left(\left(a-i\sqrt{1-a^2}\right)\phi_L(n,k)+\phi_R(n,k)\right).\end{gathered}\tag{45}$$

Local probabilities Let:

$$\begin{gathered}\Psi_{\circ}^{\dagger}\Psi_{\circ}=\rho_{\circ},\ \Psi_{\circ}'^{\dagger}\Psi_{\circ}'=\rho_{\circ}',\\ \Psi_{\circ}^{\dagger}\underline{\beta_1}\Psi_{\circ}=j_{\circ x},\ \Psi_{\circ}'^{\dagger}\underline{\beta_1}\Psi_{\circ}'=j_{\circ x}',\\ \Psi_{\circ}^{\dagger}\underline{\beta_2}\Psi_{\circ}=j_{\circ y},\ \Psi_{\circ}'^{\dagger}\underline{\beta_2}\Psi_{\circ}'=j_{\circ y}',\\ \Psi_{\circ}^{\dagger}\underline{\beta_3}\Psi_{\circ}=j_{\circ z},\ \Psi_{\circ}'^{\dagger}\underline{\beta_3}\Psi_{\circ}'=j_{\circ z}',\end{gathered}$$

$$\begin{gathered}\Psi_{*}^{\dagger}\Psi_{*}=\rho_{*},\ \Psi_{*}'^{\dagger}\Psi_{*}'=\rho_{*}',\\ \Psi_{*}^{\dagger}\underline{\beta_1}\Psi_{*}=j_{*x},\ \Psi_{*}'^{\dagger}\underline{\beta_1}\Psi_{*}'=j_{*x}',\\ \Psi_{*}^{\dagger}\underline{\beta_2}\Psi_{*}=j_{*y},\ \Psi_{*}'^{\dagger}\underline{\beta_2}\Psi_{*}'=j_{*y}',\\ \Psi_{*}^{\dagger}\underline{\beta_3}\Psi_{*}=j_{*z},\ \Psi_{*}'^{\dagger}\underline{\beta_3}\Psi_{*}'=j_{*z}',\end{gathered}$$

Because

$$\left(\Psi_\circ^\dagger\Psi_\circ\right)^2-\left(\left(\Psi_\circ^\dagger\underline{\beta_1}\Psi_\circ\right)^2+\left(\Psi_\circ^\dagger\underline{\beta_2}\Psi_\circ\right)^2+\left(\Psi_\circ^\dagger\underline{\beta_3}\Psi_\circ\right)^2\right)=$$
$$=\left(\Psi_\circ^\dagger\gamma\Psi_\circ\right)^2+\left(\Psi_\circ^\dagger\beta\Psi_\circ\right)^2,$$

$$\left(\Psi_*^\dagger\Psi_*\right)^2-\left(\left(\Psi_*^\dagger\underline{\beta_1}\Psi_*\right)^2+\left(\Psi_*^\dagger\underline{\beta_2}\Psi_*\right)^2+\left(\Psi_*^\dagger\underline{\beta_3}\Psi_*\right)^2\right)=$$
$$=\left(\Psi_*^\dagger\gamma\Psi_*\right)^2+\left(\Psi_*^\dagger\beta\Psi_*\right)^2$$

then the local densities are:

$$\rho_{\circ o}^2=\rho_\circ^2-\left(j_{\circ x}^2+j_{\circ y}^2+j_{\circ z}^2\right)=\left(\Psi_\circ^\dagger\gamma\Psi_\circ\right)^2+\left(\Psi_\circ^\dagger\beta\Psi_\circ\right)^2$$
$$\rho_{\circ o}‘^2=\rho_\circ‘^2-(j_{\circ x}‘^2+j_{\circ y}‘^2+j_{\circ z}‘^2)=$$
$$=\left(\Psi_\circ‘^\dagger\gamma\Psi_\circ‘\right)^2+\left(\Psi_\circ‘^\dagger\beta\Psi_\circ‘\right)^2$$

$$\rho_{*o}^2=\rho_*^2-\left(j_{*x}^2+j_{*y}^2+j_{*z}^2\right)=\left(\Psi_*^\dagger\gamma\Psi_*\right)^2+\left(\Psi_*^\dagger\beta\Psi_*\right)^2$$
$$\rho_{*o}‘^2=\rho_*‘^2-(j_{*x}‘^2+j_{*y}‘^2+j_{*z}‘^2)=$$
$$=\left(\Psi_*‘^\dagger\gamma\Psi_*‘\right)^2+\left(\Psi_*‘^\dagger\beta\Psi_*‘\right)^2$$

Let us design:

$$\gamma_\circ=\pi_\circ^\dagger\gamma\pi_\circ,\ \gamma_*=\pi_*^\dagger\gamma\pi_*,$$
$$\beta_\circ=\pi_\circ^\dagger\beta\pi_\circ,\ \beta_*=\pi_*^\dagger\beta\pi_*,$$

$$\gamma_\circ‘=U^{(-)\dagger}\gamma_\circ U^{(-)},\ \gamma_*‘=U^{(-)\dagger}\gamma_* U^{(-)},$$
$$\beta_\circ‘=U^{(-)\dagger}\beta_\circ U^{(-)},\ \beta_*‘=U^{(-)\dagger}\beta_* U^{(-)}.$$

Since

$$\gamma_\circ‘=a\gamma_\circ+\sqrt{1-a^2}\beta_\circ,\ \beta_\circ‘=a\beta_\circ-\sqrt{1-a^2}\gamma_\circ,$$
$$\gamma_*‘=a\gamma_*-\sqrt{1-a^2}\beta_*,\ \beta_*‘=a\beta_*+\sqrt{1-a^2}\gamma_*$$

then

$$\rho_{\circ o}^2=\rho_{\circ o}‘^2 \text{ and } \rho_{*o}^2=\rho_{*o}‘^2.$$

From (11) since:

$$\rho = \rho`,\ j_x = j_x`,\ j_y = j_y`,\ j_z = j_z`$$

then the local densities:

$$\rho_o^2 = \rho^2 - \left(j_x^2 + j_y^2 + j_z^2\right) = \rho_o`^2 = \rho`^2 - \left(j_x`^2 + j_y`^2 + j_z`^2\right).$$

Because

$$\begin{gathered}\left(\Psi^\dagger\Psi\right)^2 - \left(\left(\Psi^\dagger\underline{\beta_1}\Psi\right)^2 + \left(\Psi^\dagger\underline{\beta_2}\Psi\right)^2 + \left(\Psi^\dagger\underline{\beta_3}\Psi\right)^2\right) = \\ = \left(\Psi^\dagger\gamma\Psi\right)^2 + \left(\Psi^\dagger\beta\Psi\right)^2\end{gathered}$$

and

$$\begin{gathered}\left(\Psi`^\dagger\Psi`\right)^2 - \left(\left(\Psi`^\dagger\underline{\beta_1}\Psi`\right)^2 + \left(\Psi`^\dagger\underline{\beta_2}\Psi`\right)^2 + \left(\Psi`^\dagger\underline{\beta_3}\Psi`\right)^2\right) = \\ = \left(\begin{array}{c}\sqrt{\left(\Psi_\circ`^\dagger\gamma\Psi_\circ`\right)^2 + \left(\Psi_\circ`^\dagger\beta\Psi_\circ`\right)^2} + \\ + \sqrt{\left(\Psi_*`^\dagger\gamma\Psi_*`\right)^2 + \left(\Psi_*`^\dagger\beta\Psi_*`\right)^2}\end{array}\right)^2\end{gathered}$$

but

$$\begin{gathered}\left(\Psi`^\dagger\Psi`\right)^2 - \left(\begin{array}{c}\left(\Psi`^\dagger\underline{\beta_1}\Psi`\right)^2 + \\ + \left(\Psi`^\dagger\underline{\beta_2}\Psi`\right)^2 + \left(\Psi`^\dagger\underline{\beta_3}\Psi`\right)^2\end{array}\right) \neq \\ \neq \left(\Psi`^\dagger\gamma\Psi`\right)^2 + \left(\Psi`^\dagger\beta\Psi`\right)^2\end{gathered}$$

then

$$\begin{gathered}\rho_o = \sqrt{\left(\Psi^\dagger\gamma\Psi\right)^2 + \left(\Psi^\dagger\beta\Psi\right)^2} = \\ = \sqrt{\left(\Psi_\circ{}^\dagger\gamma\Psi_\circ\right)^2 + \left(\Psi_\circ{}^\dagger\beta\Psi_\circ\right)^2} + \\ + \sqrt{\left(\Psi_*{}^\dagger\gamma\Psi_*\right)^2 + \left(\Psi_*{}^\dagger\beta\Psi_*\right)^2} = \\ = \rho_{\circ o} + \rho_{*o}\end{gathered}$$

and

$$\begin{gathered}\rho_o` = \\ = \sqrt{\left(\Psi_\circ`^\dagger\gamma\Psi_\circ`\right)^2 + \left(\Psi_\circ`^\dagger\beta\Psi_\circ`\right)^2} + \\ + \sqrt{\left(\Psi_*`^\dagger\gamma\Psi_*`\right)^2 + \left(\Psi_*`^\dagger\beta\Psi_*`\right)^2} = \\ = \rho_{\circ o}` + \rho_{*o}`\end{gathered}$$

but

$$\rho_o{}^{\text{‘}2} \neq \left(\Psi^{\text{‘}\dagger}\gamma\Psi^{\text{‘}}\right)^2 + \left(\Psi^{\text{‘}\dagger}\beta\Psi^{\text{‘}}\right)^2.$$

Therefore ρ_o is a local probability density of a sum of two mutually exclusive events with a local densities $\rho_{\circ o}$ and ρ_{*o}.

Because:

$$\begin{array}{c}
\Psi_\circ^\dagger\gamma\Psi_\circ = \frac{1}{2}\left(1 - \frac{b}{\sqrt{1-a^2}}\right)\Psi^\dagger\gamma\Psi, \\
\Psi_\circ^\dagger\beta\Psi_\circ = \frac{1}{2}\left(1 - \frac{b}{\sqrt{1-a^2}}\right)\Psi^\dagger\beta\Psi, \\
\Psi_*^\dagger\gamma\Psi_* = \frac{1}{2}\left(1 + \frac{b}{\sqrt{1-a^2}}\right)\Psi^\dagger\gamma\Psi, \\
\Psi_*^\dagger\beta\Psi_* = \frac{1}{2}\left(1 + \frac{b}{\sqrt{1-a^2}}\right)\Psi^\dagger\beta\Psi
\end{array}$$

then ρ_o and $\rho_o{}^{\text{‘}}$ do not depend from $U^{(-)}$.

For $U^{(+)}$ (12):

$$\underline{\pi}_\circ = \frac{1}{2\sqrt{1-u^2}}\left[\begin{array}{cc}
\left(v + \sqrt{1-u^2}\right)1_4 & \left(-s + ik\right)\gamma_5 \\
\left(-ik - s\right)\gamma_5 & \left(\sqrt{1-u^2} - v\right)1_4
\end{array}\right],$$

$$\underline{\pi}_* = \frac{1}{2\sqrt{1-u^2}}\left[\begin{array}{cc}
\left(\sqrt{1-u^2} - v\right)1_4 & \left(s - ik\right)\gamma_5 \\
\left(ik + s\right)\gamma_5 & \left(v + \sqrt{1-u^2}\right)1_4
\end{array}\right].$$

Hence:

$$\begin{array}{c}
U^{(+)\dagger}\gamma U^{(+)} = u\gamma - \sqrt{1-u^2}\left(\underline{\pi}_\circ - \underline{\pi}_*\right)\beta, \\
U^{(+)\dagger}\beta U^{(+)} = u\beta + \sqrt{1-u^2}\left(\underline{\pi}_\circ - \underline{\pi}_*\right)\gamma
\end{array}$$

and all rest for $U^{(+)}$ like to $U^{(-)}$.

References

[1] G. A. Quznetsov, The Poincare group deduced from the logic properties of the information in Photon : Old Problems in Light of New Ideas, V. Dvoeglazov (ed), Nova Science Publishers, NY, 2001 or http://arXiv.org/abs/physics/9901039

[2] E.Madelung, *Die Mathematischen Hilfsmittel des Physikers.* (Springer Verlag, 1957)

[3] Gunn A. Quznetsov, The probability in the relativistic m+1 space-time, http://arXiv.org/abs/physics/9803035

[4] G. Quznetsov, The Probability Distribution to Leptons and Quarks, http://arXiv.org/abs/physics/9904024

[5] Gunn Quznetsov, The lepton, quark and hadron currents, http://arXiv.org/abs/physics/9806007.

[6] G. Quznetsov, Whence the Gauge Fields arise, http://arXiv.org/abs/physics/0006036

[7] for instance, Gordon Kane, *Modern Elementary Particle Physics.* (Addison-Wesley Publishing Company, Inc., 1987)